

Arsenic hyperaccumulating ferns: How do they survive?

June 8 2010



Researchers discover mechanisms of arsenic hyperaccumulation and tolerance using the fern *Pteris vittata*. Credit: Jo Ann Banks

Arsenic is toxic to most forms of life, and occurs naturally in soil and ground water in many regions of the world. Chronic exposure to arsenic has been linked to lung, bladder and kidney cancer, and thus there are strict limits on allowable levels of arsenic in drinking water. Chemically similar to phosphorus, arsenic forms arsenate (AsO_4^{3-}), which closely resembles phosphate (PO_4^{3-}). Arsenate interferes with many phosphate-requiring metabolic reactions, including synthesis of adenosine triphosphate (ATP), a ubiquitous and essential source of cellular energy. Thus, exposure to even low levels of arsenic can be extremely toxic.

In well-aerated soils, [arsenic](#) exists mainly as arsenate, which is taken up

by [plant roots](#) using a [phosphate](#) transporter protein. Plant tissues rapidly reduce arsenate to arsenite (AsO_3^{3-}), which is transported to the aerial portions of the plant. In aquatic environments or water-logged soils, arsenic exists primarily as arsenite. Whereas rice grains can accumulate up to 60 $\mu\text{g/g}$ arsenic, the fern *Pteris vittata* (see figure) can hyperaccumulate arsenic to levels 1000-fold greater than this. A team of researchers led by David Salt and Jo Ann Banks of Purdue University have recently isolated a gene encoding an arsenite transporter protein. This transporter allows these ferns to sequester arsenic in the vacuole, a cellular storage compartment isolated from the cytoplasm by the vacuolar membrane.

In research published this week in *The Plant Cell* and performed primarily by graduate student Emily Indriolo (now a researcher at the University of Toronto), these scientists describe how they used an arsenic-sensitive strain of yeast to isolate and characterize a gene encoding the *P. vittata* arsenite transporter. Yeast cells are arsenic-resistant because their plasma membrane contains an arsenite effluxer protein that is encoded by the *Arsenical Compound Resistance 3* (*ACR3*) gene. The researchers introduced a library of *P. vittata* [genes](#) into an arsenic-sensitive *acr3* mutant yeast strain and isolated a gene that restored arsenic resistance to this mutant. The protein encoded by this gene was then demonstrated to be very similar to the *ACR3* protein of yeast in both structure and function.

Sequence analysis showed that this fern protein contains 10 putative transmembrane domains, suggesting a cellular membrane location. Using an antibody that specifically recognizes the *ACR3* protein, they showed that *ACR3* is found in the membranes of vacuoles, but not in the plasma membrane or in endoplasmic reticulum membranes. This suggests a mechanism for arsenic tolerance in *P. vittata* tissues: arsenite that enters the cell is transported by *ACR3* into the vacuolar compartment, where it is spatially isolated from the cell cytoplasm, the site of many of the cell's

arsenic-sensitive metabolic reactions.

Furthermore, the researchers showed that *ACR3* gene expression in *P. vittata* is induced more than 30-fold in the presence of arsenite. To verify that *ACR3* is required for arsenic tolerance, the *ACR3* gene was silenced using an inhibitory mRNA. In these silenced plants, *ACR3* expression was not induced by arsenite, and arsenic significantly reduced the growth rate of these *ACR3*-deficient plants relative to unsilenced plants.

Sequence analysis showed that, although this gene is found in a wide range of organisms including bacteria, fungi, mosses and gymnosperms, it is absent in angiosperms. By studying the occurrence and function of *ACR3* in various plants, including hyperaccumulating and nonaccumulating ferns, the authors hope to provide additional insights into mechanisms of arsenic transport, tolerance, and accumulation. In addition to potential benefits for human health, this research will hopefully lead to strategies for phytoremediation of arsenic-contaminated [soil](#) and water.

More information: [www.plantcell.org/cgi/content/ ... act/tpc.109.069773v1](http://www.plantcell.org/cgi/content/act/tpc.109.069773v1)

Provided by American Society of Plant Biologists

Citation: Arsenic hyperaccumulating ferns: How do they survive? (2010, June 8) retrieved 22 September 2024 from <https://phys.org/news/2010-06-arsenic-hyperaccumulating-ferns-survive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.