

New measurements from MINOS experiment suggest a difference in a key property of neutrinos and antineutrinos

June 14 2010



Scientists know that there exist three types of neutrinos and three types of antineutrinos. Cosmological observations and laboratory-based experiments indicate that the masses of these particles must be extremely small: Each neutrino and antineutrino must weigh less than a millionth of the weight of an electron.


(PhysOrg.com) -- Scientists of the MINOS experiment at the Department of Energy's Fermi National Accelerator laboratory today announced the world's most precise measurement to date of the parameters that govern antineutrino oscillations, the back-and-forth transformations of antineutrinos from one type to another.

This result provides information about the difference in mass between different antineutrino types. The measurement showed an unexpected variance in the values for neutrinos and antineutrinos. This mass difference parameter, called Δm^2 ("delta m squared"), is smaller by approximately 40 percent for neutrinos than for antineutrinos.

However, there is still a five percent probability that Δm^2 is actually the same for neutrinos and antineutrinos. With such a level of uncertainty, MINOS physicists need more data and analysis to know for certain if the variance is real.

Neutrinos and antineutrinos behave differently in many respects, but the MINOS results, presented today at the Neutrino 2010 conference in Athens, Greece, and in a seminar at Fermilab, are the first observation of a potential fundamental difference that established physical theory could not explain.

“Everything we know up to now about neutrinos would tell you that our measured mass difference parameters should be very similar for neutrinos and antineutrinos,” said MINOS co-spokesperson Rob Plunkett. “If this result holds up, it would signal a fundamentally new property of the neutrino-antineutrino system. The implications of this difference for the physics of the universe would be profound.”



Neutrino oscillations depend on two parameters: the square of the neutrino mass difference, Δm^2 , and the mixing angle, $\sin^2 2\theta$. MINOS results (shown in black), accumulated since 2005, yield the most precise known value of Δm^2 , namely $\Delta m^2 = 0.0024 \pm 0.0001 \text{ eV}^2$.

The NUMI beam is capable of producing intense beams of either antineutrinos or neutrinos. This capability allowed the experimenters to measure the unexpected mass difference parameters. The measurement also relies on the unique characteristics of the MINOS detector, particularly its magnetic field, which allows the detector to separate the positively and negatively charged muons resulting from interactions of antineutrinos and neutrinos, respectively. MINOS scientists have also updated their measurement of the standard oscillation parameters for muon neutrinos, providing an extremely precise value of Δm^2 .

Muon antineutrinos are produced in a beam originating in Fermilab's Main Injector. The antineutrinos' extremely rare interactions with matter allow most of them to pass through the Earth unperturbed. A small number, however, interact in the MINOS detector, located 735 km away from Fermilab in Soudan, Minnesota. During their journey, which lasts 2.5 milliseconds, the particles oscillate in a process governed by a difference between their mass states.

“We do know that a difference of this size in the behavior of neutrinos and antineutrinos could not be explained by current theory,” said MINOS co-spokesperson Jenny Thomas. “While the [neutrinos](#) and antineutrinos do behave differently on their journey through the Earth, the Standard Model predicts the effect is immeasurably small in the MINOS experiment. Clearly, more antineutrino running is essential to clarify whether this effect is just due to a statistical fluctuation.”

The MINOS experiment involves more than 140 scientists, engineers, technical specialists and students from 30 institutions, including universities and national laboratories, in five countries: Brazil, Greece, Poland, the United Kingdom and the United States.

Provided by Fermilab

Citation: New measurements from MINOS experiment suggest a difference in a key property of neutrinos and antineutrinos (2010, June 14) retrieved 19 September 2024 from <https://phys.org/news/2010-06-minos-difference-key-property-neutrinos.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--