


# Researchers demonstrate highly directional Terahertz laser rays

August 8 2010

---



The metamaterial patterns are directly sculpted on the highly doped GaAs facet of the device. Artificial coloring in the figure indicates deep and shallow micron scale grooves, which have different functions. The shallow "blue" grooves efficiently couple laser output into surface electromagnetic waves on the facet and confine the waves to the facet. Credit: Courtesy of the laboratory of Federico Capasso, Harvard School of Engineering and Applied Sciences

A collaborative team of applied scientists from Harvard University and the University of Leeds have demonstrated a new terahertz (THz) semiconductor laser that emits beams with a much smaller divergence than conventional THz laser sources. The advance, published in the August 8th issue of *Nature Materials*, opens the door to a wide range of applications in terahertz science and technology. Harvard has filed a


broad patent on the invention.

The finding was spearheaded by postdoctoral fellow Nanfang Yu and Federico Capasso, Robert L. Wallace Professor of Applied Physics and Vinton Hayes Senior Research Fellow in Electrical Engineering, both of Harvard's School of Engineering and Applied Sciences (SEAS), and by a team led by Edmund Linfield at the School of Electronic and Electrical Engineering, University of Leeds.

Terahertz rays ([T-rays](#)) can penetrate efficiently through paper, clothing, plastic, and many other materials, making them ideal for detecting concealed weapons and biological agents, imaging tumors without harmful side effects, and spotting defects, such as cracks, within materials. THz radiation is also used for high-sensitivity detection of tiny concentrations of interstellar chemicals.

"Unfortunately, present THz semiconductor lasers are not suitable for many of these applications because their beam is widely divergent—similar to how light is emitted from a lamp" says Capasso. "By creating an artificial optical structure on the facet of the laser, we were able to generate highly collimated (i.e., tightly bound) rays from the device. This leads to the efficient collection and high concentration of power without the need for conventional, expensive, and bulky lenses."

Specifically, to get around the conventional limitations, the researchers sculpted an array of sub-wavelength-wide grooves, dubbed a metamaterial, directly on the facet of quantum cascade lasers. The devices emit at a frequency of 3 THz (or a wavelength of one hundred microns), in the invisible part of the spectrum known as the far-infrared.



The deep "pink" grooves form an effective grating that coherently scatters the energy of the surface waves into the far-field. Credit: Courtesy of the laboratory of Federico Capasso, Harvard School of Engineering and Applied Sciences

"Our team was able to reduce the divergence angle of the beam emerging from these [semiconductor lasers](#) dramatically, whilst maintaining the high output optical power of identical unpatterned devices," says Linfield. "This type of laser could be used by customs officials to detect illicit substances and by pharmaceutical manufacturers to check the quality of drugs being produced and stored."

The use of metamaterials, artificial materials engineered to provide properties which may not be readily available in Nature, was critical to the researchers' successful demonstration. While metamaterials have potential use in novel applications such as cloaking, negative refraction and high resolution imaging, their use in semiconductor devices has been very limited to date.

"In our case, the metamaterial serves a dual function: strongly confining the THz light emerging from the device to the laser facet and collimating the beam," explains Yu. "The ability of metamaterials to confine strongly THz waves to surfaces makes it possible to manipulate them efficiently for applications such as sensing and THz optical circuits."

Provided by Harvard University

Citation: Researchers demonstrate highly directional Terahertz laser rays (2010, August 8)  
retrieved 19 April 2024 from <https://phys.org/news/2010-08-highly-terahertz-laser-rays.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.