


CERN completes transition to lead-ion running at the Large Hadron Collider

8 November 2010



Lead-lead collisions at CMS

Four days is all it took for the LHC operations team at CERN to complete the transition from protons to lead ions in the LHC. After [extracting the final proton beam](#) of 2010 on 4 November, commissioning the lead-ion beam was underway by early afternoon. First collisions were recorded at 00:30 CET on 7 November, and stable running conditions marked the start of physics with heavy ions at 11:20 CET today.

"The speed of the transition to lead ions is a sign of the maturity of the LHC," said [CERN](#) Director General Rolf Heuer. "The machine is running like clockwork after just a few months of routine operation."

Operating the LHC with lead ions - lead atoms stripped of electrons - is completely different from operating the machine with [protons](#). From the source to collisions, operational parameters have to be re-established for the new type of beam. For lead-ions, as for protons before them, the procedure started with threading a single beam round the ring in one direction and steadily increasing the number of laps before repeating the process for the other beam. Once circulating beams had been established they could be accelerated to the full energy of 287 TeV per beam. This energy is much higher than for proton

beams because lead ions contain 82 protons. Another period of careful adjustment was needed before lining the beams up for collision, and then finally declaring that nominal data taking conditions, known at CERN as stable beams, had been established. The three experiments recording data with lead ions, ALICE, ATLAS and CMS can now look forward to continuous lead-ion running until CERN's winter technical stop begins on 6 December.

"It's been very impressive to see how well the LHC has adapted to lead [ions](#)," said Jurgen Schukraft, spokesperson of the ALICE experiment. "The ALICE detector has been optimised to record the large number of tracks that emerge from ion collisions and has handled the first collisions very well, so we are all set to explore this new opportunity at LHC."

"After a very successful proton run, we're very excited to be moving to this new phase of LHC operation," said ATLAS spokesperson Fabiola Gianotti. "The ATLAS detector has recorded first spectacular heavy-ion events, and we are eager to study them in detail."

"We designed CMS as a multi-purpose detector," said Guido Tonelli, the collaboration's spokesperson, "and it's very rewarding to see how well it's adapting to this new kind of collision. Having data collected by the same detector in proton-proton and heavy-ion modes is a powerful tool to look for unambiguous signatures of new states of matter."

Lead-ion running opens up an entirely new avenue of exploration for the LHC programme, probing matter as it would have been in the first instants of the Universe's existence. One of the main objectives for lead-ion running is to produce tiny quantities of such matter, which is known as quark-gluon plasma, and to study its evolution into the kind of matter that makes up the Universe today.

This exploration will shed further light on the properties of the strong interaction, which binds the particles called quarks, into bigger objects, such as protons and neutrons.

Following the winter technical stop, operation of the collider will start again with protons in February and physics runs will continue through 2011.

Provided by CERN

APA citation: CERN completes transition to lead-ion running at the Large Hadron Collider (2010, November 8) retrieved 4 December 2022 from <https://phys.org/news/2010-11-cern-transition-lead-ion-large-hadron.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.