


New ways to tune electrical conductivity revealed by electron interaction

January 7 2011



The energetic states of Sr₂IrO₄. The left panel shows the spectrum of all the computed electronic states. A gap in energy (y-axis) is clearly visible for electrons moving in all crystallographic directions (x-axis). The letters on the x-axis denote specific crystallographic directions. The right panel shows an isolation of the gap arising from states that form as a result of the spin-orbit interaction. Credit: Reproduced, with permission, from Ref. 1 2010 The American Physical Society

Researchers in Japan have demonstrated why the material Sr₂IrO₄, -- a transition metal oxide -- that was expected to be an electrical conductor is actually an insulator. Harnessing this material's unusual conducting properties could form the basis for novel electronic devices or superconductors.

The difference between an [electrical conductor](#) and an insulator is that [electrons](#) in the latter cannot move freely through the crystal. This is because insulators have a gap in their energy spectrum that electrons

cannot overcome. Hiroshi Watanabe, Tomonori Shirakawa and Seiji Yunoki from the RIKEN Advanced Science Institute in Wako and the Japan Science and Technology Agency have now uncovered how the electronic gap in Sr_2IrO_4 arises. Other RIKEN scientists had shown previously that the compound is an [insulator](#).

Sr_2IrO_4 is a member of the oxygen-containing compounds based on transition metals that have high atomic numbers. In these transition metals, the electrons of elements such as nickel, copper or cobalt strongly interact with each other, which results in effects such as [superconductivity](#) or magnetism.

In compounds made from the heavier transition metals, the outermost electrons circle the atoms in the so-called ‘5d electron shell’, which is relatively distant from the core. For electrons that occupy this shell there is an unusually strong interaction between their magnetic property, called spin, and the orbital motion around the atomic nucleus. The energy of this spin–orbit interaction is as large as the electron’s energy of motion or the energy arising from the electrostatic interaction between the electrons. This has dramatic consequences on their electronic properties, according to Yunoki, who led the research team. “Literally anything can happen in 5d electron systems because of the subtle balance of those three fundamental energy scales.”

How this energetic interplay modifies the electron conducting behavior in Sr_2IrO_4 became evident from the researchers’ calculations. The strong spin–orbit interaction in Sr_2IrO_4 shifts some of the electronic states to higher energies, which is sufficiently strong to create an energy gap in the electronic states.

Furthermore, the calculations reveal an intriguing connection to the family of high-temperature superconductors that have a similar gap in their electronic states. In these compounds, superconductivity is

achieved through a small addition of atoms introducing an electron surplus. The researchers are now investigating the possibility that this could also be the case here. “It would have an enormous impact if one can make Sr_2IrO_4 superconducting,” says Yunoki. “We hope that our theoretical calculations will be of help to experimentalists.”

More information:

- Watanabe, H., et al. Microscopic study of a spin-orbit-induced Mott insulator in Ir oxides. *Physical Review Letters* 105, 216410 (2010).
- Kim, B., et al. Phase-sensitive observation of a spin-orbital Mott state in Sr_2IrO_4 . *Science* 323, 1329–1332 (2009).

Provided by RIKEN

Citation: New ways to tune electrical conductivity revealed by electron interaction (2011, January 7) retrieved 25 April 2024 from

<https://phys.org/news/2011-01-ways-tune-electrical-revealed-electron.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.