


# Mathematical model could help predict and prevent future extinctions

January 25 2011



When an ecosystem is healthy, food, territory and other resources are balanced among species, as depicted in the top left diagram. If a key species or network component drastically decreases, an entire system may be disrupted as shown on the left. Aiding the troubled species or component, the approach illustrated in the top right image, may further strain the connections in the network, perpetuating the imbalance. A newly developed mathematical model suggests some extinctions or network failures can be avoided and the network can be rebalanced by suppressing, rather than enhancing, one or more members of the network, bottom right. Credit: Zina Deretsky, National Science Foundation

In an effort to better understand the dynamics of complex networks, scientists have developed a mathematical model to describe interactions within ecological food webs. This research, performed by Northwestern University physics professor Adilson Motter and his student, Sagar Sahasrabudhe, is published in the January 25 issue of *Nature Communications*. The work illustrates how human intervention may

effectively aid species conservation efforts.

"Our study provides a theoretical basis for management efforts that would aim to mitigate [extinction](#) cascades in [food web](#) networks. There is evidence that a significant fraction of all extinctions are caused not by a primary perturbation but instead by the propagation of a cascade," said Motter.

Extinction cascades are often observed following the loss of a key species within an ecosystem. As the system changes to compensate for the loss, availability of food, territory and other resources to each of the remaining members can fluctuate wildly, creating a boom-or-bust environment that can lead to even more extinctions. According to the study, more than 70 percent of these extinctions are preventable, assuming that the system can be brought into balance using only available resources--no new factors may be introduced.

Motter explained further, "We find that extinction cascades can often be mitigated by suppressing--rather than enhancing--the populations of specific species. In numerous cases, it is predicted that even the proactive removal of a species that would otherwise be extinct by a cascade can prevent the extinction of other species."

The finding may seem counterintuitive to conservationists because the compensatory actions seem to inflict further damage to the system. However, when the entire ecosystem is considered, the effect is beneficial. This news holds promise for those charged with maintaining Earth's biodiversity and natural resources--the health of which can counteract many of the causes of climate change, and some man-made disasters such as the Gulf of Mexico oil spill.

The dodo bird, *Raphus cucullatus*, is one example of extinction due to human activity. The dodo was a large, flightless bird that became extinct

in the 1600s. It is likely that a combination of factors including hunting, loss of habitat, and perhaps even a flash flood, stressed the ecosystem on the island of Mauritius, home of the dodo. Some researchers think that human introduction of non-native species, such as dogs, pigs, cats and rats to the island, is what ultimately lead to the demise of the dodo.

In any case, in the future, it may be possible to avoid extinction of some species in stressed ecosystems by applying the new method of analysis developed by Motter.

The goal of this project, funded by the National Science Foundation's Division of Mathematical Sciences, is to develop mathematical methods to study dynamical processes in [complex networks](#). Although the specific application mentioned here may be useful in management of ecosystems, the mathematical foundation underlying the analysis is much more universal. The broad concept is innovative in the area of complex networks because it concludes that large-scale failures can be avoided by focusing on preventing the waves of failure that follow the initial event.

This approach could be used to stabilize a wide array of complex networks. It can apply to biochemical networks in order to slow or stop progression of diseases caused by variations inside individual cells. It can also be used to manage technological networks such as the smart grid to prevent blackouts. It can even apply to regulation of complicated financial networks by identifying key factors in the early stages of a financial downturn, which, when met with human intervention, could potentially save billions of dollars.

The world is a complicated place that gets even trickier when trying to mathematically explain a complex network, especially when the network evolves within an environment that is itself changing. But, Motter says his [mathematical model](#) is promising for the study of changing environments.

"Uncertainty itself is not a problem," he quipped. "The problem comes when you cannot estimate uncertainty."

Provided by National Science Foundation

Citation: Mathematical model could help predict and prevent future extinctions (2011, January 25) retrieved 20 April 2024 from <https://phys.org/news/2011-01-mathematical-future-extinctions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.