

Oldest fossils of large seaweeds, worm-like animals tell story of ancient oxygen

February 23 2011



This cluster of fossils from the Lantian area was preserved and unearthed in very high densities. Credit: Zhe Chen and Xunlai Yuan

Almost 600 million years ago, before the rapid evolution of life forms known as the Cambrian explosion, a community of seaweeds and worm-like animals lived in a quiet deep-water niche near what is now Lantian, a small village in south China.

Then they simply died, leaving some 3,000 nearly pristine fossils preserved between beds of black shale deposited in oxygen-free and unbreathable waters.

Scientists from the Chinese Academy of Sciences, Virginia Tech in the United States and Northwest University in Xi'an, China report the discovery of the fossils in the Feb. 17 issue of the journal *Nature*.

In addition to ancient versions of algae and worms, the Lantian biota--named for its location--included microfossils with complex and puzzling structures.

In all, scientists have identified some 15 species at the site.

The fossils suggest that structural diversification of macroscopic eukaryotes--the earliest versions of organisms with complex cell structures--may have occurred only tens of millions of years after the [Snowball Earth](#) event that ended 635 million years ago.

Snowball Earth proposes that the Earth's surface became almost, or completely, frozen at least once during the planet's history.

The presence of macroscopic eukaryotes in the highly organic-rich black shale suggests that, despite the overall oxygen-free conditions, brief [oxygenation](#) of the oceans did come and go, according to H. Richard Lane, program director in the National Science Foundation (NSF)'s Division of Earth Sciences, which funded the research.

"So there are two questions," says Shuhai Xiao, a geobiologist at Virginia Tech. "Why did this community evolve when, and where, it did?"


"It is clearly different in terms of the number of species compared to biota preserved in older rocks. There are more species here, and they are more complex and larger than what evolved before."

The rocks were formed shortly after the largest ice age ever, says Xiao, when much of the global ocean was frozen.

By 635 million years ago, the snowball Earth event ended and the oceans were clear of ice. Perhaps, Xiao says, "that prepared the ground for the evolution of complex eukaryotes."

The team examined the black shale rocks because, although they were laid down in less than optimal waters for oxygen-dependent organisms, "they are known to be able to preserve fossils very well," says Shuhai.

"In most cases, dead organisms were washed in and preserved in black shales. In this case, we discovered fossils that were preserved in pristine condition--some [seaweeds](#) still rooted--where they had lived."



This seaweed holdfast is a 600 million-year-old fossil from the Lantian Formation in China. Credit: Zhe Chen and Xunlai Yuan

The conclusion that the environment would have been poisonous is derived from geochemical data, "but the bedding surfaces where these fossils were found represent moments of geologic time during which oxygen was available and conditions were favorable," says Xiao.

"They are very brief moments to a geologist, but long enough for the oxygen-demanding organisms to colonize the Lantian basin and capture the rare opportunities."

The research team suggests that the Lantian basin was largely without oxygen, but was punctuated by brief oxic episodes that were populated by complex new life forms.

Those life forms were subsequently killed and preserved when the oxygen disappeared.

"Such brief intervals need high-resolution sampling for geochemical analysis to capture the dynamic and complex nature of oxygen history in the Ediacaran Period," says lead paper author Xunlai Yuan. The Ediacaran Period is the last geological period immediately preceding the Cambrian Period.

Proving that hypothesis awaits further study.

The rocks in the study region are deposited in layered beds. The nature of the rock changes subtly, and there are finer and finer layers that can be recognized within each bed.

"We will need to sample each layer to see whether there is any difference in oxygen contents between layers with fossils and those without," says co-author Chuanming Zhou.

Provided by National Science Foundation

Citation: Oldest fossils of large seaweeds, worm-like animals tell story of ancient oxygen (2011, February 23) retrieved 15 May 2024 from <https://phys.org/news/2011-02-oldest-fossils-large-seaweeds-worm-like.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.