

# Scientists make quantum breakthrough

20 April 2011



Artist's impression of the speckle pattern created by a multimode light beam (top, red), and the image measured in this experiment created by a multimode beam of atoms (top, blue). Source: Dr Tim Wetherell, ANU.

(PhysOrg.com) -- Scientists have demonstrated for the first time that atoms can be guided in a laser beam and possess the same properties as light guided in an optical communications fiber.

The researchers' work has implications for future quantum devices that require smoothly-guided matter waves, such as atom interferometers which need to sensitively measure the earth's gravitational field for geo-exploration. Their paper is published today in *Nature Communications*.

"In an optical fibre, many modes of light can be conducted simultaneously, and they can interfere to produce a speckled pattern of light," said team member Professor Ken Baldwin from the ARC Centre of Excellence for Quantum-Atom Optics at ANU.

"We have shown that when [atoms](#) in a vacuum chamber are guided inside a laser light beam, they too can create a speckle pattern - an image of which we have captured for the first time".

The team trapped a cloud of cold helium atoms at the focus of an intense [laser beam](#) pointed downwards at the imaging system, and then gradually turned down the laser intensity until the speckled image appeared. The work was done with PhD students Sean Hodgman and Andrew Manning.

"We then made the atoms even colder," says team leader Dr Andrew Truscott, "until they behaved more like waves than particles, forming a single quantum wave called a Bose-Einstein condensate (BEC). When the BEC was loaded into the guide, the speckle pattern disappeared, showing that just one mode was being transmitted - the single quantum wave."

The physicists demonstrated that by measuring the arrival time of the atoms on the imaging system, they were able to distinguish between the multimode (speckled image) guiding, and the single-mode (smooth image) guiding.

"Measurements for the multi-mode beam showed the atoms arriving in groups as a result of their interference - so-called atom bunching," said team member Dr Robert Dall. "However, the BEC represents just a single quantum mode with no interference, so when we guided the BEC - we saw no bunching."

The guiding behaviour agreed with a theoretical model developed by team member Mattias Johnsson. "We have shown that atoms can be guided in a laser beam of light, with the same properties as light guided in an optical fibre made of glass," said Dr Johnsson.

**More information:** Observation of atomic speckle and Hanbury Brown - Twiss correlations in guided matter waves, *Nature Communications* 2, Article number: 291 [doi:10.1038/ncomms1292](https://doi.org/10.1038/ncomms1292)

## Abstract

Speckle patterns produced by multiple independent light sources are a manifestation of the coherence

of the light field. Second-order correlations exhibited in phenomena such as photon bunching, termed the Hanbury Brown - Twiss effect, are a measure of quantum coherence. Here we observe for the first time atomic speckle produced by atoms transmitted through an optical waveguide, and link this to second-order correlations of the atomic arrival times. We show that multimode matter-wave guiding, which is directly analogous to multimode light guiding in optical fibres, produces a speckled transverse intensity pattern and atom bunching, whereas single-mode guiding of atoms that are output-coupled from a Bose - Einstein condensate yields a smooth intensity profile and a second-order correlation value of unity. Both first- and second-order coherence are important for applications requiring a fully coherent atomic source, such as squeezed-atom interferometry.

Provided by Australian National University

APA citation: Scientists make quantum breakthrough (2011, April 20) retrieved 2 March 2021 from <https://phys.org/news/2011-04-scientists-quantum-breakthrough.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*