

Scientists discover animal-like urea cycle in tiny diatoms in the ocean

May 11 2011



Micrograph of the fusiform morphotype of the diatom *Phaeodactylum tricornutum*. Credit: Alessandra De Martino, Ecole Normale Supérieure, Paris

Scientists have discovered that marine diatoms, tiny phytoplankton abundant in the sea, have an animal-like urea cycle, and that this cycle enables the diatoms to efficiently use carbon and nitrogen from their environment.

The researchers, from the J. Craig Venter Institute (JCVI) and other institutions, published their findings in this week's issue of the journal *Nature*.

The team, led by lead author Andrew Allen from JCVI and co-author Chris Bowler, Institute of Biology, Ecole Normale Supérieure, Paris, believes that the cycle could be a reason for the domination of diatoms

in marine environments, especially after upwelling events--the upward movement of nutrient rich waters from the deep ocean to the surface.


In response to ocean upwelling, diatoms are able to quickly recover from prolonged periods of nutrient deprivation and rapidly proliferate.

"This study provides fascinating insights into how diatoms have evolved to become the dominant primary producers in many ocean regions," says David Garrison, program director in the National Science Foundation's (NSF) Division of Ocean Sciences, which funded the research along with NSF's Division of Molecular and Cellular Biosciences.

Diatoms have unique cell walls made of silica. They are key organisms for understanding the environmental health of marine ecosystems, and are responsible for much of the carbon and oxygen production in the ocean.

Diatom photosynthesis in ocean environments is also responsible for about one fifth of the oxygen in the atmosphere.

In previous research, Allen, Bowler and colleagues sequenced the genome of the first pennate diatom, *Phaeodactylum tricorutum*.



Diatoms from Puget Sound, Washington; diatoms are a critical part of the

marine environment Credit: Adrian Marchetti, University of Washington, and Andrew Allen, JCVI

In that research, they developed new methods for determining the origin of diatom genes. They also looked at nutrient metabolism in diatoms, beginning with iron metabolism.

Building on that work, Allen and colleagues explored the evolutionary history of diatoms, specifically *P. tricornutum*, and cellular mechanisms for nutrient utilization in the environment, leading to the finding that diatoms have a functional urea cycle.

This was a stunning discovery, says Allen, because it was thought that the urea cycle originated with the metazoan (animal) branch of life.

There it has played an important role in facilitating a wide range of physiological innovations in vertebrates.

For example, urea synthesis enables rapid control of minerals and salts in the blood in animals such as sharks, skates, rays and bony fish, and ammonia detoxification associated with water retention in amphibians and mammals.

The latter was likely a prerequisite for life on land, and subsequently enabled the biochemical pathways necessary for processing a high-protein diet.

Allen and others have now shown that the urea cycle originated hundreds of millions of years before the appearance of metazoans.

The team used RNA interference techniques to partially silence a key

changes the way we view diatoms relative to animals and plants."

The work also suggests that diatoms have followed a fundamentally different evolutionary path from plants--the dominant oxygen producers in terrestrial environments, green algae, and other closely related organisms.

Rather, prior to evolutionary acquisition of photosynthetic machinery, the ancestors of diatoms were possibly more closely related to the ancestors of animals than to plants.

This relatedness has resulted in diatoms and animals sharing some similar biochemical pathways such as the urea cycle.

Although it appears that animals and diatoms ultimately use the urea cycle for different purposes, they are evolutionarily linked in a way that animals and plants are not.

More information: Andrew E. Allen, Christopher L. Dupont, Miroslav Oborník, Aleš Horák, Adriano Nunes-Nesi,, John P. McCrow, Hong Zheng, Hanhua Hu, Alisdair R. Fernie, Chris Bowler, Exosymbiont-derived urea cycle used for intracellular carbon and nitrogen recovery in diatoms, *Nature*, May 12th 2011, [DOI: 10.1038/nature10074](https://doi.org/10.1038/nature10074)

Provided by National Science Foundation

Citation: Scientists discover animal-like urea cycle in tiny diatoms in the ocean (2011, May 11) retrieved 25 April 2024 from <https://phys.org/news/2011-05-urea-anabolic-steroid-diatoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.