


Physicists find charge separation in a molecule consisting of two identical atoms

25 November 2011



Electron density in a homonuclear molecule which behaves like a polar dimer molecule. The size of this molecule is more than 1000 times larger than usual molecules. Image: Science/AAAS

Physicists from the University of Stuttgart show the first experimental proof of a molecule consisting of two identical atoms that exhibits a permanent electric dipole moment. This observation contradicts the classical opinion described in many physics and chemistry textbooks. The work was published in the journal *Science* yesterday.

A dipolar molecule forms as a result of a [charge separation](#) between the negative charged [electron cloud](#) and the positive core, creating a permanent [electric dipole moment](#). Usually this charge separation originates in different attraction of the cores of different elements onto the negative charged [electrons](#). Due to symmetry reasons homonuclear molecules, consisting only of atoms of the same element, therefore could not possess dipole moments.

However, the dipolar molecules that were

discovered by the group of Prof. Tilman Pfau at the 5th Institute of Physics at the University of Stuttgart do consist of two atoms of the element rubidium. The necessary [asymmetry](#) arises as a result of different electronically excited states of the two alike atoms. Generally this excitation will be exchanged between the atoms and the asymmetry will be lifted. Here this exchange is suppressed by the huge size of the molecule, which is about 1000 times larger than an [oxygen molecule](#) and reaches sizes of viruses. Therefore the probability to exchange the [excitation](#) between the two atoms is so small that it would statistically only happen once in the lifetime of the universe. Consequently, these homonuclear molecules possess a dipole moment. A permanent dipole moment additionally requires an orientation of the molecular axis. Due to their size the molecules rotate so slowly that the dipole moment does not average out from the viewpoint of an observer.

Physicists from the University of Stuttgart succeeded in experimentally detecting the dipole moment. They measured the energy shift of the molecule in an electric field by [laser spectroscopy](#) in an ultra cold atomic cloud. The same group caused worldwide a stir when they created these weakly bound Rydberg molecules for the first time in 2009. The molecules consist of two identical atoms whereof one is excited to a highly excited state, a so-called Rydberg state. The unusual binding mechanism relies on scattering of the highly excited Rydberg electron of the second atom. So far theoretical descriptions of this binding mechanism did not predict a dipole moment. However, the scattering of the Rydberg electron of the bound atom changes the probability distribution of the electron. This breaks the otherwise spherical symmetry and creates a dipole moment. In collaboration with theoretical physicists from the Max-Planck-Institute for the Physics of Complex Systems in Dresden and from the Harvard-Smithsonian Center for Astrophysics in Cambridge, USA, a new theoretical treatment was developed

that confirms the observation of a dipole moment.

The proof of a permanent dipole moment in a homonuclear molecule not only improves the understanding of polar molecules. Ultra cold polar molecules are also promising to study and control chemical reactions of single molecules.

More information: A Homonuclear Molecule with a Permanent Electric Dipole Moment, *Science* 25 November 2011: Vol. 334 no. 6059 pp. 1110-1114.
[DOI: 10.1126/science.1211255](https://doi.org/10.1126/science.1211255)

ABSTRACT

Permanent electric dipole moments in molecules require a breaking of parity symmetry. Conventionally, this symmetry breaking relies on the presence of heteronuclear constituents. We report the observation of a permanent electric dipole moment in a homonuclear molecule in which the binding is based on asymmetric electronic excitation between the atoms. These exotic molecules consist of a ground-state rubidium (Rb) atom bound inside a second Rb atom electronically excited to a high-lying Rydberg state. Detailed calculations predict appreciable dipole moments on the order of 1 Debye, in excellent agreement with the observations.

Provided by University of Stuttgart

APA citation: Physicists find charge separation in a molecule consisting of two identical atoms (2011, November 25) retrieved 23 May 2019 from <https://phys.org/news/2011-11-physicists-molecule-identical-atoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.