


Vampire star reveals its secrets

7 December 2011



These super-sharp images of the unusual vampire double star system SS Leporis were created from observations made with the VLT Interferometer at ESO's Paranal Observatory using the PIONIER instrument. The system consists of a red giant star orbiting a hotter companion. The remarkable image sharpness — 50 times sharper than those from the NASA/ESA Hubble Space Telescope — not only allows the stars to be clearly separated and their orbital motion followed, but also allowed the size of the red giant to be measured more accurately than ever before. The system consists of a red giant star orbiting a hotter companion. Note that the stars have been artificially coloured to match their known temperatures. Credit: ESO/PIONIER/IPAG

(PhysOrg.com) -- Astronomers have obtained the best images ever of a star that has lost most of its material to a vampire companion. By combining the light captured by four telescopes at ESO's Paranal Observatory they created a virtual telescope 130 metres across with vision 50 times sharper than the NASA/ESA Hubble Space Telescope. Surprisingly, the new results show that the transfer of mass from one star to the other in this double system is gentler than expected.

"We can now combine light from four VLT telescopes and create super-sharp images much more quickly than before," says Nicolas Blind (IPAG, Grenoble, France), who is the lead author on the paper presenting the results, "The images are so sharp that we can not only watch the [stars](#) orbiting around each other, but also measure the size of the larger of the two stars."

The astronomers observed the unusual system SS Leporis in the constellation of Lepus (The Hare), which contains two stars that circle around each other in 260 days. The stars are separated by only a little more than the distance between the Sun and the Earth, while the largest and coolest of the two stars extends to one quarter of this distance -- corresponding roughly to the orbit of Mercury. Because of this closeness, the hot companion has already cannibalised about half of the mass of the larger star.

"We knew that this double star was unusual, and that material was flowing from one star to the other," says co-author Henri Boffin, from ESO. "What we found, however, is that the way in which the mass transfer most likely took place is completely different from previous models of the process. The 'bite' of the vampire star is very gentle but highly effective."

The new observations are sharp enough to show that the giant star is smaller than previously thought, making it much more difficult to explain how the red giant lost matter to its companion. The astronomers now think that, rather than streaming from one star to the other, the matter must be expelled from the giant star as a stellar wind and captured by the hotter companion.

"These observations have demonstrated the new snapshot imaging capability of the Very Large [Telescope](#) Interferometer. They pave the way for many further fascinating studies of interacting double stars," concludes co-author Jean-Philippe Berger.

More information: This research was presented in a paper, "An incisive look at the symbiotic star SS Leporis - Milli-arcsecond imaging with PIONIER/VLTI", by N. Blind et al. in press in the journal *Astronomy & Astrophysics*.

Provided by ESO

APA citation: Vampire star reveals its secrets (2011, December 7) retrieved 16 June 2019 from <https://phys.org/news/2011-12-vampire-star-reveals-secrets.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.