



calendar year of compilation, according to the 2011 State of Observed Species report released Jan. 18, 2012, by the International Institute for Species Exploration at Arizona State University. More at <http://species.asu.edu>. Credit: International Institute for Species Exploration/Arizona State University

"It has been speculated, for example, that marine microbial species alone could number 20 million," he said.

With those staggering numbers as a backdrop, statistics, or "species bites," from the latest report note that:


- Almost 24 percent of the new vascular plant species discovered in 2009 were in the monocot order *Asparagales*, which includes orchids, hyacinths, irises, daffodils, amaryllis, allium, aloe and, of course, asparagus.
- Year to year, the largest order of newly discovered [insects](#) is the beetles, and, 2009 was no exception. Overall, 3,485 new beetle species (*Coleoptera*) were officially described including rove beetles (568), ground beetles (421), long-horned beetles (369), leaf beetles (356) and scarabs (288).
- Only 41 new living mammal species were officially described in 2009 and of those, 83 percent were either bats (44 percent) or rodents (39 percent).
- Almost 90 percent (133) of the new living amphibian species described in 2009 were frogs.
- There was almost five times more fossil bird species (34) newly described in 2009 than living birds (seven).
- Typical of most years, the largest number of new fish species was in the order *Perciformes* and 29 percent of those were in the families *Gobiidae* (22) and *Cichlidae* (11). Gobies include some of the tiniest fish on Earth, and the cichlids include some of the most popular aquarium fish, including the angelfish and damselfish.
- Of the 626 newly described living crustacean species, 224 (31.8 percent) were in the order *Decapoda*, which includes

crayfish, crabs, lobsters, prawns and shrimp.

- The *Colubridae* is the largest family of snakes and in 2009, almost 65 percent of the newly described living snakes were colubrids. In addition to 31 new snakes, new reptile species (living) included 38 lizards, 29 geckos, 12 iguanas, five chameleons and two turtles.
- More than 13 percent of the new fungus species (living) described in 2009 were gilled mushrooms in the order *Agaricales* (178). Of the mushrooms, more than one-fifth (21.3 percent) were in the family *Marasmiaceae*, which includes shiitake mushrooms.

In addition to the living species discovered during 2009, there were 1,905 fossil species, with insects and spiders accounting for 25.6 percent.

"As the number of species increases, so too does our understanding of the biosphere," said Wheeler, a professor in the School of Sustainability and a Senior Sustainability Scientist in the Global Institute of Sustainability at ASU. "It is through knowledge of the unique attributes of species that we illuminate the origin and evolutionary history of life on our planet. As we find out where species live and how they interact, we increase our ability to understand the function of ecosystems and make effective, fact-based decisions regarding conservation."



This chart illustrates a decade of the discovery of species newly known to science, which is detailed in the Retro SOS (State of Observed Species) report released Jan. 18, 2012, by the International Institute for Species Exploration at Arizona State University. It visually shows there were 176,311 newly discovered and officially described from 2000-2009. More at <http://species.asu.edu>. Credit: International Institute for Species Exploration/Arizona State University

This is the fourth year for the annual State of Observed Species report compiled by the International Institute for Species Exploration. In addition to the 2011 report, the institute is also releasing a Retro SOS - a decade of species discovery in review - 2000-2009. The Retro SOS notes that from 2000 through 2009, there were 176,311 newly discovered species.

Provided by Arizona State University

"It is particularly instructive to understand the tempo and patterns of discovery in recent years," said Wheeler, adding, "Given this data, it is interesting to ponder underlying causes of trends."

The "obvious lesson" from compiling this data, according to Wheeler, is that all nomenclatural acts, including descriptions of new species, must be mandatorily registered going forward. "In the animal world it takes about two years to mine the international literature for evidence of newly named species. The current lack of registration requirements simply compounds the problem of an already massive backlog," he said.

The report notes there are increasing calls for more aggressive and visionary approaches to mapping the species of the biosphere. "The adaptation of cyberinfrastructure to eliminate bottlenecks in the practice of taxonomy has created an opportunity to vastly accelerate species exploration," said Wheeler, who uses the SOS report and the annual naming of the top 10 new species each May, as ways to draw attention to this mission.

The SOS report and the Retro SOS are filled with statistics and charts, including a colorful word cloud. Sara Pennak, assistant director for partnerships and public outreach at the institute, prepared the data synthesis and analysis for the reports, which are available online at <http://species.asu.edu>.

Partners in this effort include: Algae Base, MycoBank, International Journal of Systematic and Evolutionary Microbiology, World Register of Marine [Species](#) (WoRMS), Thomson Reuters Zoological Record, International Plant Names Index, UniProt and Taxatoy.

APA citation: Insects top latest inventory of newly discovered species (2012, January 18) retrieved 6 December 2022 from <https://phys.org/news/2012-01-insects-latest-newly-species.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*