

Radiation detected 400 miles off Japanese coast

21 February 2012, By BRIAN SKOLOFF and MALCOLM RITTER , Associated Press



Members of the media wearing protective suits and masks report, escorted by an official of Nuclear and Industrial Safety Agency, second left, at Tokyo Electric Power Co.'s tsunami-crippled Fukushima Dai-ichi nuclear power plant in Fukushima prefecture, northeastern Japan, Monday, Feb. 20, 2012. Japan next month marks one year since the March 11 tsunami and earthquake, which triggered the worst nuclear accident since Chernobyl in 1986. (AP Photo/Issei Kato, Pool)

(AP) -- Radioactive contamination from the Fukushima power plant disaster has been detected as far as almost 400 miles off Japan in the Pacific Ocean, with water showing readings of up to 1,000 times more than prior levels, scientists reported Tuesday.

But those results for the substance cesium-137 are far below the levels that are generally considered harmful, either to [marine animals](#) or people who eat seafood, said Ken Buesseler of the Woods Hole Oceanographic Institution.

He spoke Tuesday in [Salt Lake City](#) at the annual Ocean Sciences Meeting, attended by more than 4,000 researchers this week.

The results are for [water samples](#) taken in June, about three months after the power plant disaster, Buesseler said. In addition to thousands of water

samples, researchers also sampled fish and plankton and found cesium-137 levels well below the legal health limit.

"We're not over the hump" yet in terms of [radioactive contamination](#) of the ocean because of continued leakage from the plant, Buesseler said in an interview before Tuesday's talk. He was chief scientist for the cruise that collected the data.

The ship sampled water from about 20 miles to about 400 miles off the coast east of the Fukushima plant. Concentrations of cesium-137 throughout that range were 10 to 1,000 times normal, but they were about one-tenth the levels generally considered harmful, Buesseler said.

Cesium-137 wasn't the only [radioactive substance](#) released from the plant, but it's of particular concern because of its long persistence in the environment. Its half-life is 30 years.

The highest readings last June were not always from locations closest to the Fukushima plant, Buesseler said. That's because swirling [ocean currents](#) formed concentrations of the material, he said.

Most of the cesium-137 detected during the voyage probably entered the ocean from water discharges, rather than atmospheric fallout, he added.

Hartmut Nies of the [International Atomic Energy Agency](#) said Buesseler's findings were not surprising, given the vastness of the ocean and its ability to absorb and dilute materials.

"This is what we predicted," Nies said after Buesseler presented his research.

Nies said the water's cesium-137 concentration has been so diluted that just 20 miles offshore, "if it was not seawater, you could drink it without any problems."

"This is good news," he said, adding that scientists expect levels to continue to decrease over time.

"We still don't have a full picture," Nies said, "but we can expect the situation will not become worse."

©2012 The Associated Press. All rights reserved.

This material may not be published, broadcast, rewritten or redistributed.

APA citation: Radiation detected 400 miles off Japanese coast (2012, February 21) retrieved 20 October 2019 from <https://phys.org/news/2012-02-miles-japanese-coast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.