

# Immortal worms defy aging

February 27 2012

---



Polycelis felina, a freshwater planarian. Image: Eduard Sola, Wikipedia.

Researchers from The University of Nottingham have demonstrated how a species of flatworm overcomes the ageing process to be potentially immortal.

The discovery, published in the [Proceedings of the National Academy of Sciences](#), may shed light on the possibilities of alleviating ageing and age-related characteristics in [human cells](#).

Planarian worms have amazed scientists with their apparently limitless ability to regenerate. Researchers have been studying their ability to replace aged or damaged tissues and cells in a bid to understand the mechanisms underlying their [longevity](#).

Dr Aziz Aboobaker from the University's School of Biology, said:  
"We've been studying two types of planarian worms; those that reproduce sexually, like us, and those that reproduce asexually, simply dividing in two. Both appear to regenerate indefinitely by growing new muscles, skin, guts and even entire brains over and over again.

"Usually when stem cells divide — to heal wounds, or during reproduction or for growth — they start to show signs of ageing. This means that the stem cells are no longer able to divide and so become less able to replace exhausted specialised cells in the tissues of our bodies. Our ageing skin is perhaps the most visible example of this effect. Planarian worms and their stem cells are somehow able to avoid the [ageing process](#) and to keep their cells dividing."

One of the events associated with ageing cells is related to telomere length. In order to grow and function normally, cells in our bodies must keep dividing to replace cells that are worn out or damaged. During this division process, copies of the genetic material must pass on to the next generation of cells. The genetic information inside cells is arranged in twisted strands of DNA called chromosomes. At the end of these strands is a protective 'cap' called a telomere. Telomeres have been likened to the protective end of a shoelace which stops strands from fraying or sticking to other strands.

Each time a cell divides the protective telomere 'cap' gets shorter. When they get too short, the cell loses its ability to renew and divide. In an immortal animal we would therefore expect cells to be able to maintain telomere length indefinitely so that they can continue to replicate. Dr Aboobaker predicted that planarian worms actively maintain the ends of their chromosomes in adult stem cells, leading to theoretical immortality.

Dr Thomas Tan made some exciting discoveries for this paper as part of his PhD. He performed a series of challenging experiments to explain

the worm's immortality. In collaboration with the rest of the team, he also went some way to understanding the clever molecular trick that enabled cells to go on dividing indefinitely without suffering from shortened chromosome ends.

Previous work, leading to the award of the 2009 Nobel Prize for Physiology or Medicine, had shown that telomeres could be maintained by the activity of an enzyme called telomerase. In most sexually reproducing organisms the enzyme is most active only during early development. So as we age, telomeres start to reduce in length.

This project identified a possible planarian version of the gene coding for this enzyme and turned down its activity. This resulted in reduced telomere length and proved it was the right gene. They were then able to confidently measure its activity and resulting telomere length and found that asexual worms dramatically increase the activity of this gene when they regenerate, allowing [stem cells](#) to maintain their telomeres as they divide to replace missing tissues.

Dr Tan pointed out the importance of the interdisciplinary expertise: "It was serendipitous to be sandwiched between Professor Edward Louis's yeast genetics lab and the Children's [Brain](#) Tumour Research Centre, both University of Nottingham research centres with expertise in telomere biology. Aziz and Ed kept demanding clearer proof and I feel we have been able to give a very satisfying answer."

However, what puzzled the team is that sexually reproducing planarian worms do not appear to maintain telomere length in the same way. The difference they observed between asexual and sexual animals was surprising, given that they both appear to have an indefinite regenerative capacity. The team believe that sexually reproductive worms will eventually show effects of telomere shortening, or that they are able to use another [mechanism](#) to maintain telomeres that would not involve the

telomerase enzyme.

Dr Aboobaker concluded: "Asexual planarian [worms](#) demonstrate the potential to maintain telomere length during regeneration. Our data satisfy one of the predictions about what it would take for an animal to be potentially immortal and that it is possible for this scenario to evolve. The next goals for us are to understand the mechanisms in more detail and to understand more about how you evolve an immortal animal."

Professor Douglas Kell, BBSRC Chief Executive, said: "This exciting research contributes significantly to our fundamental understanding of some of the processes involved in ageing, and builds strong foundations for improving health and potentially longevity in other organisms, including humans."

Provided by University of Nottingham

Citation: Immortal worms defy aging (2012, February 27) retrieved 26 April 2024 from <https://phys.org/news/2012-02-immortal-worms-defy-aging.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.