

# Removal of invasive tree improves health of American Samoa forests

March 26 2012

---

Removal of the Tamaligi tree (*Falcataria moluccana*), an invasive and destructive non-native tree on Tutuila Island, American Samoa greatly improves the health of its diverse native forests, according to a recently published study appearing in the journal *Biological Invasions*.

A team of scientists from the USDA [Forest](#) Service's Pacific Southwest Research Station–Hilo and the National Park of American Samoa (NPSA) investigated how Tamaligi [trees](#) affected the composition, biomass, and soil nitrogen in forests within or adjacent to the NPSA. The team sampled both forests with Tamaligi, and forests where Tamaligi had been present, but subsequently removed.

Native to the Moluccas, New Guinea, New Britain, and the Solomon Islands, Tamaligi was introduced to Tutuila Island in the early 1900s. Aided by its nitrogen-fixing capacity, Tamaligi—among the world's fastest growing trees—thrives on a variety of soil types, including degraded sites and acidic or nutrient-poor soils. By 2000, 35 percent of Tutuila Island was infested with the invasive tree, prompting NPSA to begin control of this species.

Since 2001, there has been an aggressive field campaign to eliminate the Tamaligi tree within the boundaries of the NPSA and surrounding areas. Invasive species control in general, and control of the Tamaligi tree in particular, is considered a high priority as Samoa's forests are important constituents of the Pacific Ocean's Polynesia/Micronesia biodiversity hotspot. These islands support some of the most intact native ecosystems

of any Pacific Island group, including shoreline, or littoral, communities, wetlands, and lowland, montane and cloud rainforests.

Findings from the current study demonstrated the strong influence of Tamaligi on the structure, composition, and functioning of American Samoa native forests. That Tamaligi-invaded forests exhibited equivalent levels of total biomass, but significantly lower biomass of native tree species, suggests that this invasion replaces, rather than augments, the biomass of native trees in these forests. But once Tamaligi trees are controlled, the native trees quickly recover and shade out any future Tamaligi regeneration.

"The effort to eliminate Tamaligi populations from the NPSA is a great and inspiring example of successful control of an invasive species," says Dr. R. Flint Hughes, a PSW research ecologist who led the study.

"Funding effectively supported the control efforts. Widespread public support was garnered through outreach with local village leadership, employment of villagers, and use of the media. And even though Tamaligi is a daunting invasive, it can be controlled, and when it is, many of the native Samoan tree species recover and persist."

**More information:** To read the full article, go to:  
[treesearch.fs.fed.us/pubs/40018](https://treesearch.fs.fed.us/pubs/40018)

Provided by USDA Forest Service

Citation: Removal of invasive tree improves health of American Samoa forests (2012, March 26) retrieved 20 September 2024 from  
<https://phys.org/news/2012-03-invasive-tree-health-american-samoa.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.