

Fertilizer use responsible for increase in nitrous oxide in atmosphere

2 April 2012

University of California, Berkeley, chemists have found a smoking gun proving that increased fertilizer use over the past 50 years is responsible for a dramatic rise in atmospheric nitrous oxide, which is a major greenhouse gas contributing to global climate change.

Climate scientists have assumed that the cause of the increased [nitrous oxide](#) was nitrogen-based fertilizer, which stimulates [microbes](#) in the soil to convert nitrogen to nitrous oxide at a faster rate than normal.

The new study, reported in the April issue of the journal [Nature Geoscience](#), uses nitrogen isotope data to identify the unmistakable fingerprint of fertilizer use in archived [air samples](#) from Antarctica and Tasmania.

"Our study is the first to show empirically from the data at hand alone that the nitrogen isotope ratio in the atmosphere and how it has changed over time is a fingerprint of fertilizer use," said study leader Kristie Boering, a UC Berkeley professor of chemistry and of earth and [planetary science](#).

"We are not vilifying fertilizer. We can't just stop using fertilizer," she added. "But we hope this study will contribute to changes in fertilizer use and [agricultural practices](#) that will help to mitigate the release of nitrous oxide into the atmosphere."

Since the year 1750, nitrous oxide levels have risen 20 percent - from below 270 parts per billion (ppb) to more than 320 ppb. After carbon dioxide and methane, nitrous oxide (N₂O) is the most potent greenhouse gas, trapping heat and contributing to global warming. It also destroys stratospheric ozone, which protects the planet from harmful ultraviolet rays.

Not surprisingly, a steep ramp-up in atmospheric nitrous oxide coincided with the green revolution that increased dramatically in the 1960s, when

inexpensive, synthetic fertilizer and other developments boosted food production worldwide, feeding a burgeoning global population.

Tracking the origin of nitrous oxide in the atmosphere, however, is difficult because a molecule from a fertilized field looks identical to one from a natural forest or the ocean if you only measure total concentration. But a quirk of microbial metabolism affects the isotope ratio of the nitrogen the N₂O microbes give off, producing a telltale fingerprint that can be detected with sensitive techniques.

Archived air from Cape Grim

Boering and her colleagues, including former UC Berkeley graduate students Sunyoung Park and Phillip Croteau, obtained air samples from Antarctic ice, called firn air, dating from 1940 to 2005, and from an atmospheric monitoring station at Cape Grim, Tasmania, which has archived air back to 1978.

Analysis of N₂O levels in the Cape Grim air samples revealed a seasonal cycle, which has been known before. But isotopic measurements by a very sensitive isotope ratio mass spectrometer also displayed a seasonal cycle, which had not been observed before. At Cape Grim, the isotopes show that the seasonal cycle is due both to the circulation of air returning from the stratosphere, where N₂O is destroyed after an average lifetime of 120 years, and to seasonal changes in the ocean, most likely upwelling that releases more N₂O at some times of year than at others.

"The fact that the isotopic composition of N₂O shows a coherent signal in space and time is exciting, because now you have a way to differentiate agricultural N₂O from natural ocean N₂O from Amazon forest emissions from N₂O returning from the stratosphere," Boering said. "In addition, you also now have a way to check

whether your international neighbors are abiding by agreements they've made to mitigate N₂O emissions. It is a tool that, ultimately, we can use to verify whether N₂O emissions by agriculture or biofuel production are in line with what they say they are."

Changes in fertilizer use can reduce N₂O emissions

Limiting nitrous oxide emissions could be part of a first step toward reducing all greenhouse gases and lessening global warming, Boering said, especially since immediately reducing global carbon dioxide emissions is proving difficult from a political standpoint. In particular, reducing nitrous oxide emissions can initially offset more than its fair share of [greenhouse gas](#) emissions overall, since N₂O traps heat at a different wavelength than CO₂ and clogs a "window" that allows Earth to cool off independent of CO₂ levels.

"On a pound for pound basis, it is really worthwhile to figure how to limit our emissions of N₂O and methane," she said. "Limiting N₂O emissions can buy us a little more time in figuring out how to reduce CO₂ emissions."

One approach, for example, is to time fertilizer application to avoid rain, because wet and happy soil microbes can produce sudden bursts of nitrous oxide. Changes in the way fields are tilled, when they are fertilized and how much is used can reduce N₂O production.

Boering's studies, which involve analyzing the isotopic fingerprints of nitrous oxide from different sources, could help farmers determine which strategies are most effective. It could also help assess the potential negative impacts of growing crops for biofuels, since some feedstocks may require fertilizer that will generate N₂O that offsets their carbon neutrality.

"This new evidence of the budget of nitrous oxide allows us to better predict its future changes - and therefore its impacts on climate and stratospheric ozone depletion - for different scenarios of fertilizer use in support of rising populations and increased production for bio-energy," said coauthor David

Etheridge of the Centre for Australian Weather and Climate Research in Aspendale, Victoria.

Finding the fingerprint of fertilized microbes

Boering was able to trace the source of N₂O because bacteria in a nitrogen rich environment, such as a freshly fertilized field, prefer to incorporate nitrogen-14 (¹⁴N), the most common isotope, instead of nitrogen-15 (¹⁵N).

"Microbes on a spa weekend can afford to discriminate against nitrogen-15, so the fingerprint of N₂O from a fertilized field is a greater proportion of nitrogen-14," Boering said. "Our study is the first to show empirically from the data at hand alone that the nitrogen isotope ratio in the atmosphere and how it has changed over time is a fingerprint of fertilizer use."

Just as telling is the isotope ratio of the central nitrogen atom in the N-N-O molecule. By measuring the nitrogen isotope ratio overall, the isotope ratio in the central nitrogen atom, and contrasting these with the oxygen-18/oxygen-16 isotope ratio, which has not changed over the past 65 years, they were able to paint a consistent picture pointing at fertilizer as the major source of increased atmospheric N₂O .

The isotope ratios also revealed that [fertilizer](#) use has caused a shift in the way soil microbes produce N₂O. The relative output of bacteria that produce N₂O by nitrification grew from 13 to 23 percent worldwide, while the relative output of bacteria that produce N₂O by denitrification - typically in the absence of oxygen - dropped from 87 to 77 percent. Although the numbers themselves are uncertain, these are the first numerical estimates of these global trends over time, made possible by the unique archived air dataset of this study.

Provided by University of California - Berkeley

APA citation: Fertilizer use responsible for increase in nitrous oxide in atmosphere (2012, April 2)
retrieved 20 September 2019 from <https://phys.org/news/2012-04-fertilizer-responsible-nitrous-oxide-atmosphere.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.