


# A new optical microscopy approach opens the door to better observations in molecular biology

17 May 2012



Super-resolution optical reconstruction of HIV morphology. Average distribution of the integrase enzyme as visualized by FIAsH-PALM (top panel). The high resolution of this technique (~30nm) allows to recover the characteristic conical shape of the capsid. By contrast, conventional microscopy (resolution ~200-300 nm) cannot reveal details of this structure (bottom panel). Credit: Pasteur Institut

Researchers from the Institut Pasteur and CNRS have set up a new optical microscopy approach that combines two recent imaging techniques in order to visualize molecular assemblies without affecting their biological functions, at a resolution 10 times better than that of traditional microscopes.

Using this approach, they were able to observe the AIDS virus and its capsids (containing the HIV genome) within cells at a scale of 30 nanometres, for the first time with light. This newly developed approach represents a significant advance in molecular biology, opening the door to less invasive and more precise analyses of pathogenic microorganisms present in human host cells. This study is already published in the Electronic Edition of *PNAS*.

It has always been necessary for researchers to visualise [pathogenic microorganisms](#) in their host cell's environment, in order to define the host-pathogen interactions contributing to viral infections. Optical microscopy, combined with fluorescent labels (such as GFP proteins and antibodies coupled with synthetic fluorophores), allows to showcase the specific structures of cells, including proteins. However, this approach is limited by its low resolving power, which only helps distinguish cellular and [molecular structures](#) at a scale of 200-300 nanometres (nm). Most [pathogenic viruses](#) are of smaller sizes. Consequently, it is essential to resort to more precise imaging techniques, in order to better understand and define the [internal structure](#) of such viruses.

A study coordinated by Dr. Christophe Zimmer, in collaboration with Dr. Nathalie Arhel within the lab headed by Pr Pierre Charneau, shows that the association of two recent imaging techniques helps obtain unique images of molecular assemblies of HIV-1 capsids, with a resolution around 10 times better than that of traditional microscopes. This new approach, which uses super-resolution imaging and FIAsH labeling, does not affect the virus' ability to self-replicate. It represents a major step forward in molecular biology studies, enabling the visualisation of microbial complexes at a scale of

30 nm without affecting their function.

The newly developed approach combines super-resolution PALM imaging and fluorescent FIAsh labeling. PALM imaging relies on the acquisition of thousands of low-resolution images, each of which showing only a few fluorescent molecules. The molecular positions are then calculated with high accuracy by computer programs and compiled into a single high-resolution image. FIAsh labeling involves the insertion of a 6-amino-acid peptide into the protein of interest. The binding of the FIAsh fluorophore to the peptide generates a fluorescent signal, thereby enabling the visualization of the protein. For the first time, researchers have combined these two methods in order to obtain high-resolution images of molecular structures in either fixed or living cells.

This new method has helped researchers visualise the [AIDS Virus](#) and localize its capsids in human cells, at a scale of 30 nm. Capsids are conical structures which contain the [HIV genome](#). These structures must dismantle in order for the viral genome to integrate itself into the host cell's genome. However, the timing of this disassembly has long been debated. According to a prevailing view, capsids disassemble right after infection of the [host cell](#) and, therefore, do not play an important role in the intracellular transport of the virus to the host cell's nucleus. However, the results obtained by the researchers of the Institut Pasteur and CNRS indicate that numerous capsids remain unaltered until entry of the virus into the nucleus, confirming and strengthening earlier studies based on electron microscopy. Hence, capsids could play a more important role than commonly assumed in the replication cycle of HIV.

The development of a new [optical microscopy](#) approach by the researchers of the Institut Pasteur and CNRS offers unique perspectives for [molecular biology](#). This new imaging technique could become a key tool in the study of numerous microbial complexes and their interactions with host cells at the molecular level. This non-invasive technique allows to observe proteins without destroying or altering their biological functions. Moreover, this technique could eventually enable the analysis of microorganisms with single-nanometre accuracy,

thereby ensuring a transition from microscopy to "nanoscopy". Consequently, the next steps are the sharing of this new approach with the scientific community, its further development and its application to the study of other pathogenic microorganisms.

**More information:** Super resolution imaging of HIV in infected cells with FIAsh-PALM - online Electronic Edition of *PNAS* - May 14, 2012. Mickaël Lelek et al.

Provided by CNRS

APA citation: A new optical microscopy approach opens the door to better observations in molecular biology (2012, May 17) retrieved 20 June 2021 from <https://phys.org/news/2012-05-optical-microscopy-approach-door-molecular.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*