

Diseased trees new source of climate gas

7 August 2012

Diseased trees in forests may be a significant new source of methane that causes climate change, according to researchers at the Yale School of Forestry & Environmental Studies in *Geophysical Research Letters*.

Sixty trees sampled at Yale Myers [Forest](#) in northeastern Connecticut contained concentrations of [methane](#) that were as high as 80,000 times ambient levels. Normal air concentrations are less than 2 parts per million, but the Yale researchers found average levels of 15,000 parts per million inside trees.

"These are flammable concentrations," said Kristofer Covey, the study's lead author and a Ph.D. candidate at Yale. "Because the conditions thought to be driving this process are common throughout the world's forests, we believe we have found a globally significant new source of this potent greenhouse gas."

The estimated emission rate from an upland site at the Yale forest is roughly equivalent to burning 40 gallons of gasoline per hectare of forest per year. It also has a global warming potential equivalent to 18 percent of the carbon being sequestered by these forests, reducing their climate benefit of carbon sequestration by nearly one-fifth.

"If we extrapolate these findings to forests globally, the methane produced in trees represents 10 percent of global emissions," said Xuhui Lee, a co-author of the study and Sara Shallenberger Brown Professor of Meteorology at Yale. "We didn't know this pathway existed."

The trees producing methane are older-between 80 and 100 years old-and diseased. Although outwardly healthy, they are being hollowed out by a common fungal infection that slowly eats through the trunk, creating conditions favorable to methane-producing microorganisms called methanogens.

"No one until now has linked the idea that fungal rot of timber trees, a production problem in

commercial forestry, might also present a problem for greenhouse gas and climate change mitigation," said Mark Bradford, a co-author and Assistant Professor of Terrestrial Ecosystem Ecology at F&ES.

Red maple, an abundant species in North America, had the highest methane concentrations, but other common species, including oak, birch and pine were also producers of the gas. The rate of methane emissions was 3.1 times higher in the summer, suggesting that higher temperatures may lead to increasing levels of forest methane that, in turn, lead to ever-higher temperatures.

"These findings suggest decay in living [trees](#) is important to biogeochemists and atmospheric scientists seeking to understand global greenhouse gas budgets and associated [climate change](#)," said Covey.

More information: "Elevated Methane Concentrations in Trees of an Upland Forest," www.agu.org/pubs/crossref/pip/2012GL052361.shtml

Provided by Yale University

APA citation: Diseased trees new source of climate gas (2012, August 7) retrieved 11 April 2021 from <https://phys.org/news/2012-08-diseased-trees-source-climate-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.