

All about dust

August 28 2012



The Cygnus-X star formation region as seen in the infrared with the Herschel Space Telescope. Most of this image is produced by warm dust heated by ultraviolet light from the new stars. A new paper presents a new, more accurate technique for deriving the temperature and properties of the dust from such images. Credit: NASA/ESA/Herschel, M.Hennemann & F. Motte

(Phys.org)—The space between stars is not empty—it contains copious amounts of gas and dust. Astronomers estimate that about 5-10% of the total mass of our Milky Way galaxy is contained in the interstellar gas and dust, and about 1% is in the form of tiny dust grains made predominantly of silicates, rather like fine sand. Some grains are also composed of carbon and other elements.

[Interstellar dust](#) plays many key roles in astronomy. Dust absorbs visible light, and so determines in large part what astronomers can see in the visible. At the same time, dust emits at [infrared wavelengths](#), and by

virtue of this process of absorption and subsequent re-emission of light dust controls much of the [energy balance](#) in the interstellar medium. Dust is also essential to the chemistry that takes place in the interstellar medium because it provides [gas molecules](#) with a surface on which to react with other molecules. Dust grains themselves contain a large fraction of many important elements in the universe like silicon, carbon, or iron. Not least, at an early stage in the evolution of [new stars](#) the dust around them can coagulate into clumps—a step towards the formation of planets.

Warm dust in space—that is, dust heated by ultraviolet from new stars to the relatively "warm" temperatures of 20 to 60 kelvin—has recently become the subject of intense study thanks to the Herschel [Space Observatory](#). This telescope has instruments designed to image this faint, extended material. Modeling the emission precisely, however, has turned out to be a challenge. One issue is that the color of the emitted radiation depends not only on the temperature but also on the dust grains' size, shape and composition. So far most astronomers have tried to fit their observations to best fits using a simple algorithm that minimizes the cumulative difference between the observed and predicted values at all different wavelengths.

CfA astronomers Brandon Kelly and Alyssa Goodman, with four colleagues, have made significant improvements to the technique of determining dust temperatures and properties from infrared observations. Rather than relying on traditional minimization techniques, they simultaneously fit a set of data points and model predictions (using a so-called Bayesian algorithm), and extract a set of most probable dust values. Their new techniques, when applied to Herschel data, obtain some significant results. Rather than the traditional finding that dust temperature is anti-correlated with its properties like size (that is, hot dust grains are smaller), they find the opposite: a weak correlation between temperature and these properties. The new approach is still

being applied, with more results expected, but the work seems to promise a more precise and reliable way to measure the properties of dust, a key component of the interstellar medium.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: All about dust (2012, August 28) retrieved 25 April 2024 from <https://phys.org/news/2012-08-all-about-dust.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.