

# Finding undetonated bombs on the sea floor

13 November 2012



CSIRO researcher, Keith Leslie, prepares the magnetic sensor for field trials.

(Phys.org)—CSIRO has developed a sensor to detect undetonated explosives on the sea floor. It is based on technology used to find mineral deposits underground.

The sensor was developed as part of a project with US Government agency, the Strategic Environmental Research and Development Program (SERDP) and US-based research organisation Sky Research.

The method for finding undetonated underwater explosives is very similar to that used to detect underground [mineral deposits](#), says CSIRO electrical engineer Dr Keith Leslie.

"Our highly sensitive sensor – the high temperature superconducting tensor [gradiometer](#) – delivers

significantly more information about the [target](#)'s magnetic field than conventional sensors used for this type of detection," he said.

"It provides data on the location, characterisation and magnetic qualities of a target – whether it is a gold deposit or an explosive."

Over 10 million acres of coastal waters are contaminated by undetonated explosives, according to SERDP. Typically these small explosives rust and corrode at sea, making them even more dangerous.

"The marine environment is difficult to sample due to electrical currents produced by waves, which affect underwater magnetic fields," Dr Leslie said.

"In [mineral exploration](#), near [surface deposits](#) are being exhausted, leading our search for minerals deeper underground, where targets are more difficult to detect with traditional surface and airborne measurements."

The CSIRO sensor can provide valuable geological information that discriminates between prospective and non-prospective areas or targets. It avoids unnecessary drilling and minimises the risk of overlooking valuable mineral deposits.

"Our sensor has a critical advantage for small targets such as undetonated explosives, where only one or two measurements may be near the target," Dr Leslie said.

"In mineral exploration, a string of measurements of the gradients of the magnetic field down a [drill hole](#) can determine the direction to the target."

Eventually the technology may renew exploration efforts at abandoned sites where drilling programs were based on insufficient or inaccurate information. It also has the potential to help clear landmines.

The sensor has been proved in a stationary

laboratory environment. Trials have been conducted to prove it in motion, in preparation for anticipated underwater trials.

Provided by CSIRO

APA citation: Finding undetonated bombs on the sea floor (2012, November 13) retrieved 15 September 2019 from <https://phys.org/news/2012-11-undetonated-sea-floor.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*