

Our galaxy's 'geysers' are towers of power

2 January 2013



This shows the "geysers" (in blue) shooting out of the Milky Way. Credit: Optical image – A. Mellinger, U. Central Michigan; radio image – E. Carretti, CSIRO; radio data – S-PASS team; composition – E. Bresser, CSIRO

"Monster" outflows of charged particles from the centre of our Galaxy, stretching more than halfway across the sky, have been detected and mapped with CSIRO's 64-m Parkes radio telescope. Corresponding to the "Fermi Bubbles" found in 2010, the outflows were detected by astronomers from Australia, the USA, Italy and The Netherlands. The finding is reported in today's issue of *Nature*.

"These outflows contain an extraordinary amount of energy—about a million times the energy of an [exploding star](#)," said the research team's leader, CSIRO's Dr Ettore Carretti.

But the outflows pose no danger to Earth or the Solar System.

The speed of the outflow is supersonic, about 1000 kilometres a second. "That's fast, even for astronomers," Dr Carretti said. "They are not coming in our direction, but go up and down from the Galactic Plane. We are 30,000 light-years away from the Galactic Centre, in the Plane. They

are no danger to us."

From top to bottom the outflows extend 50,000 light-years [five hundred thousand million million kilometres] out of the [Galactic Plane](#). That's equal to half the diameter of our Galaxy (which is 100,000 light-years—a million million million kilometres—across).

Seen from Earth, the outflows stretch about two-thirds across the sky from horizon to horizon. The outflows correspond to a "haze" of microwave emission previously spotted by the WMAP and Planck space telescopes and regions of [gamma-ray emission](#) detected with NASA's Fermi [space telescope](#) in 2010, which were dubbed the "Fermi Bubbles".

The WMAP, Planck and Fermi observations did not provide enough evidence to indicate definitively the source of the radiation they detected, but the new Parkes observations do.

"The options were a quasar-like outburst from the black hole at the Galactic Centre, or star-power—the hot winds from [young stars](#), and exploding stars," said team member Dr Gianni Bernardi of the Harvard-Smithsonian Center for Astrophysics, in Cambridge, Massachusetts. "Our observations tell us it's star-power."

In fact, the outflows appear to have been driven by many generations of stars forming and exploding in the Galactic Centre over the last hundred million years.

The key to determining this was to measure the outflows' magnetic fields.

"We did this by measuring a key property of the radio waves from the outflows—their polarisation," said team member Dr Roland Crocker of the Max-Planck-Institut fuer Kernphysik in Heidelberg, Germany, and the Australian National University.

The new observations also help to answer one of

astronomers' big questions about our Galaxy: how it generates and maintains its magnetic field. "The outflow from the [Galactic Centre](#) is carrying off not just gas and high-energy electrons, but also strong magnetic fields," said team member Dr Marijke Haverkorn of Radboud University Nijmegen in The Netherlands.

"We suspect this must play a big part in generating the Galaxy's overall magnetic field."

Provided by CSIRO

APA citation: Our galaxy's 'geysers' are towers of power (2013, January 2) retrieved 26 September 2020 from <https://phys.org/news/2013-01-galaxy-geysers-towers-power.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.