


Violent Nabro eruption shown to pierce stratosphere

February 8 2013



A natural-color image of the Nabro eruption captured June 13, 2011, by the NASA Moderate Resolution Imaging Spectroradiometer (MODIS) shows the volcanic ash plume extending from the Eritrea-Ethiopia border, west-north-westerly, into Sudan. Nabro is a stratovolcano in the northeast African nation of Eritrea and is one of many volcanic caldera complexes in the northeasternmost part of the East African Rift valley region. Credit: NASA Earth Observatory

(Phys.org)—Researchers from the U.S. Naval Research Laboratory Ocean and Atmospheric Science and Technology Directorate report conclusive evidence that volcanic gases and aerosols from the explosive 2011 eruption of the East Africa Nabro Volcano breached the stratosphere, unassisted by the seasonal Asian monsoonal flow.


On June 12, 2011, the Nabro Volcano explosively erupted, forcing sulfur dioxide (SO₂) gas and volcanic aerosols high into the atmosphere. Prior analyses had suggested the Nabro plume reached the stratosphere only after being transported through a tropospheric meteorological phenomenon associated with the [Asian Monsoon](#) circulation.

"There is overwhelming satellite evidence that the Nabro [eruption](#) caused a sizable and direct volcanic impact on the stratosphere," said Mike Fromm, NRL research meteorologist. "Robust satellite data indisputably indicate [volcanic gases](#) and particles penetrated the stratosphere during the initial eruption and a second eruption three days later."

Thermal infrared (IR) imagery captured by the NASA Moderate Resolution Imaging Spectroradiometer (MODIS) satellite two hours after the eruption provides overwhelming evidence of a deep convective umbrella cloud with locally [cold temperatures](#) in a peripheral arc and warmer temperatures near the cloud center, a signature of violent convection associated with breaches in the [lower stratosphere](#).

To characterize the Nabro plume, NRL scientists use three instruments from NASA's 'A-Train' constellation—a series of quasi-simultaneous nadir and limb viewing measurement satellites that orbit Earth one behind the other. MODIS, onboard the satellite Aqua, acquires spectral data on global dynamics and processes occurring on land, in oceans, and in the atmosphere. The Ozone Monitoring Instrument (OMI), onboard the Aura satellite, distinguishes between aerosol types such as smoke,

dust, and sulfates, and measures data to derive ozone. The Microwave Limb Sounder (MLS), also onboard Aura, provides measurements of atmospheric composition, temperature, humidity, and cloud ice. Data taken from the OMI shortly after the eruption indicate scattering volcanic aerosols such as liquid sulfate droplets and SO₂. Observations from MODIS IR cloud-top temperatures revealed that an optically opaque volcanic cloud covers the same area as the OMI AI plume with a minimum IR brightness temperature of minus 78 degrees Celsius, a lower stratospheric value.



MODIS IR cloud-top temperature on June 13, 2011 reveals that the optically opaque volcanic cloud covers the same area as the OMI AI plume. The minimum IR brightness temperature of this cloud is -78°C (dark red) which according to the environmental temperature profile suggests an opaque cloud as high as 16 kilometers. UV backscatter sensors such as OMI are sensitive cannot 'see through' optically thick clouds. Hence the signal of scattering aerosols sulfur dioxide (SO₂) is originating from altitudes above this high cloud top. The

opaque cloud surface also guarantees that no information from the troposphere below is obtainable. Credit: U.S. Naval Research Laboratory

"Ultraviolet backscatter sensors such as OMI are sensitive and cannot 'see through' optically thick clouds, indicating the signal of scattering aerosols and sulfur dioxide originated from altitudes above this high cloud top," says Fromm. "The opaque cloud surface also guarantees that no information from the troposphere below is obtainable."

The day after the eruption, height-resolved SO₂ data retrieved from the MLS also exhibited strong stratospheric SO₂ enhancements to an altitude of nearly 63,000 feet, or approximately 19 kilometers. Hence, the SO₂ profile data corroborate the high injection heights suggested by the combined OMI/MODIS imagery. Additionally, a back trajectory launched from a June 19 MLS SO₂ observation shows that the stratospheric SO₂ plume moves from the Mideast and Africa through mid-latitude Asia to the Pacific Ocean, never encountering the Asian Monsoon—a progression consistent with the region's recorded prevailing winds.

These data convincingly show a classic Plinian characterization of the Nabro June 13 eruption defined by volcanic aerosols and other gases being injected into the stratosphere.

Provided by Naval Research Laboratory

Citation: Violent Nabro eruption shown to pierce stratosphere (2013, February 8) retrieved 22 September 2024 from <https://phys.org/news/2013-02-violent-nabro-eruption-shown-pierce.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.