

# New aplodontid rodent found from the late Oligocene of Northern Junggar Basin, China

March 22 2013




Fig.1. Upper dentition of *Proansomys durensis*. Credit: BI Shundong

Mountain beavers of the subfamily Ansominae are small-sized aplodontid rodents, characterized by a bucket-handle shaped ectoloph on their upper cheek teeth. Although fossils have been discovered from the Oligocene and Miocene of North America and Eurasia, the evolutionary

origins and biogeography of the group are still poorly known because of relatively few records and poor representation of some taxa.

Paleontologists from Institute of Vertebrate Paleontology and Paleoanthropology (IVPP), Chinese Academy of Sciences, American Museum of Natural History and Indiana University of Pennsylvania, described a new genus and species of aplodontid rodent, *Proansomys durensis*, from the late Oligocene of the northern Junggar Basin of China. These specimens represent the earliest record of the subfamily in Asia and provide new information on the early history of the Ansomylinae, as recently reported in *PLoS ONE* 8(1).

The majority of specimens were obtained by surface collecting and screen washing from the Tieersihabahe faunal assemblage Zone I of the Tieersihabahe Formation during the 1998, 1999, 2000, 2002, and 2004 expeditions, and some materials were from Saerduoyila, approximately 50 km northwest of Tieersihabahe section in the Halamagai area. All specimens are isolated brachydont teeth. The age of the fauna is biostratigraphically and paleomagnetically correlative to the late Oligocene, around 24.4-24.15 million years ago.




Fig.2. Lower dentition of *Proansomys durensis*. Credit: BI Shundong

The new genus is referred to as *Ansomyinae* because the ectoloph on the upper cheek teeth, although not fully crested, has attained the same characteristic bucket-handle-shaped configuration as other members of the subfamily. It represents the earliest record of the subfamily yet discovered in Asia and is more plesiomorphic than species of the genus *Ansomys* in having a partly crested ectoloph, a lower degree of lophodonty, and less complex tooth basins (lacking accessory lophules). *Proansomys* has transitional features between *Prosciurus* and *Ansomys*, suggesting that the *Ansomyinae* derived from a group of aplodontids related to *Prosciurus*, as did other advanced aplodontid rodents. This provides new light on the paleobiogeography of the *Ansomyinae*.

*Proansomys durensis* is assigned to Ansomyinae because the ectoloph of upper cheek teeth, although not fully crested, has attained the bucket-handle shaped configuration that is the most important diagnostic character of the subfamily as its name implies. However, it is distinct from all other known species of *Ansomys* by its significantly larger size, a partly crested ectoloph, the absence of accessory lophules, lower degree of lophodonty, and the straight protoloph with crest-like protoconule on most upper molars.

"Cladistic analysis shows the new species is more primitive than species of *Ansomys* in having a partly crested ectoloph on upper cheek teeth, less complex dental pattern without the accessory lophules, a lesser degree of lophodonty, metaconid on m1 lingually prominent, main cusps of lower teeth not anteroposteriorly compressed, and an incomplete hypolophid on p4. Therefore, we consider this justifies its status as a new genus", said study author Dr. BI Shundong from the IVPP.

There are two alternate hypotheses for the paleobiogeography of the *Ansomys*. Some proposed a European origin and a series of migrations between Europe, Asia, and North America, while some proposed an origin in North America and a single migration to Eurasia in the early Miocene. "The recognition of the lineage (*Prosciurus-Proansomys-Ansomys*) in our study provides new insights into the early history of the ansomyines. Although the earliest record of ansomyines was *A. cyanotephrus* from the early late Oligocene of North America, *Proansomys* is the most primitive member of the subfamily and appears derived from *Prosciurus*-like forms. Given that *Prosciurus* was common in the early Oligocene in Mongolia, the ansomyines may have originated in Asia some time prior to the late [Oligocene](#), then dispersed throughout Eurasia and North America", said BI Shundong, "However, more complete fossil discoveries will be necessary to make reliable inferences about the ansomyine paleobiogeographical relations".

This work was mainly supported by the [Chinese Academy of Sciences](#) (CAS) Key Project, the National Nature Science Foundation of China, the CAS 100-Talent Project, and the CAS Fossil Excavation and Preparation Fund.

**More information:** <http://%20http://www.plosone.org/article/info:doi/10.1371/journal.pone.0052625>

Provided by Institute of Vertebrate Paleontology and Paleoanthropology

Citation: New aplodontid rodent found from the late Oligocene of Northern Junggar Basin, China (2013, March 22) retrieved 22 May 2024 from <https://phys.org/news/2013-03-aplodontid-rodent-late-oligocene-northern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.