

# Lonely bees make better guests

18 June 2013, by Alex Peel



Bumblebee coated in pollen.

Solitary bees are twice as likely to pollinate the flowers they visit as their more sociable counterparts, according to a new study.

It says an oilseed rape flower visited by a solitary bee stands a 71 per cent chance of receiving pollen from the body of the bee. Should a [bumblebee](#) or honeybee stop by, those chances drop by half.

The research, published in *Agriculture, Ecosystems and Environment*, underlines the vital role that wild bees play in providing back-up pollination services to our struggling [honeybee colonies](#), and the growing need to protect them.

'Underlying all of this is a lack of [honeybees](#),' says Dr Ben Woodcock from NERC's Centre for [Ecology](#) & Hydrology, who led the study. 'There simply aren't enough honeybee colonies in the UK to effectively pollinate all of our crops.'

'They're not going to do it alone; they need support from wild bee populations.'

The team spent two months on oilseed rape farms in Wiltshire, closely monitoring the bees as they meandered from flower to flower.

[Solitary bees](#) spent longer at [flowers](#), and were more likely to have pollen available on their bodies

than honeybees or bumblebees.

They and bumblebees also made more than twice as much contact with the stigma, the pollen-collecting part of the flower, as honeybees.

Honeybees are one of the UK's most important pollinators, often occurring in high numbers near managed colonies. But the number of managed colonies has fallen by more than half in 20 years, and a survey out last week says more than a third of UK hives have failed to survive the cold, wet winter.



Solitary bee

Those declines have had a dramatic effect on honeybees' ability to pollinate crops. In 1984, they were capable of satisfying up to 70 per cent of [pollen](#) demand for UK crops, but a study in 2011 said that figure had dropped to just 34 per cent.

'In the UK, honeybees near arable crops are often managed with the goal of producing honey' explains Woodcock. 'They're not often managed with the goal of pollinating arable crops such as oilseed rape or field beans'

'For practical reasons hives are often clustered together in fairly small areas of the landscape, so you can get a scenario where wild bees make an

important contribution to pollination away from those clusters.'

'If we're going to ensure that our crops are pollinated, we're going to need a robust wild bee population.'

'They need species-rich grasslands, diverse field margins, and foraging resources that are available outside of the crop-flowering season.'

**More information:** Woodcock, B., Crop flower visitation by honeybees, bumblebees and solitary bees: Behavioural differences and diversity responses to landscape, *Agriculture, Ecosystems and Environment*, 2013. [DOI: 10.1016/j.agee.2013.03.005](https://doi.org/10.1016/j.agee.2013.03.005)

*This story is republished courtesy of [Planet Earth online](http://www.planetearthonline.org), a free, companion website to the award-winning magazine Planet Earth published and funded by the Natural Environment Research Council (NERC).*

Provided by PlanetEarth Online

APA citation: Lonely bees make better guests (2013, June 18) retrieved 25 November 2020 from <https://phys.org/news/2013-06-lonely-bees-guests.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*