

# Not the end of the world: Why Earth's greatest mass extinction was the making of modern mammals

28 August 2013



Skeleton of the cynodont *Galesaurus planiceps*. Credit: Roger Smith, Iziko Museums of South Africa Social History / Natural History / Art Collections

The ancient closest relatives of mammals – the cynodont therapsids - not only survived the greatest mass extinction of all time, 252 million years ago, but thrived in the aftermath, according to new research published today.

The first mammals arose in the Triassic period, more than 225 million years ago. These early fur balls include small shrew-like animals such as *Morganucodon* from England, *Megazostrodon* from South Africa and *Bienotherium* from China.

They had differentiated teeth (incisors, canines, molars) and large brains and were probably warm-blooded and covered in fur – all characteristics that stand them apart from their reptile ancestors, and which contribute to their huge success today.

However, new research suggests that this array of unique features arose gradually over a long span of time, and that the first mammals may have

arisen as a result of the end-Permian [mass extinction](#) - which wiped out 90 per cent of [marine organisms](#) and 70 per cent of [terrestrial species](#).

The research was conducted by the University of Lincoln, UK, the National Museum in Bloemfontein, South Africa, and the University of Bristol, UK, and has been published today in *Proceedings of the Royal Society B*.

Lead author Dr Marcello Ruta, evolutionary palaeobiologist from the University of Lincoln's School of Life Sciences, said: "Mass extinctions are seen as entirely negative. However, in this case, cynodont therapsids, which included a very small number of species before the extinction, really took off afterwards and were able to adapt to fill many different niches in the Triassic - from carnivores to herbivores."

Co-author Dr Jennifer Botha-Brink of the National Museum in Bloemfontein, South Africa, said: "During the Triassic, the cynodonts split into two groups, the cynognathians and the probainognathians. The first were mainly plant-eaters, the second mainly flesh-eaters and the two groups seemed to rise and fall at random - first one expanding, and then the other. In the end, the probainognathians became the most diverse and most varied in adaptations, and they gave rise to the first mammals some 25 million years after the mass extinction."

Co-author Professor Michael Benton added: "We saw that when a major group, such as cynodonts, diversifies, it is the body shape or range of adaptations that expands first. The diversity, or number of species, rises after all the morphologies available to the group have been tried out."

The researchers concluded that cynodont diversity

rose steadily during the recovery of life following the mass extinction, with their range of form rising rapidly at first before hitting a plateau. This suggests there is no particular difference in morphological diversity between the very first [mammals](#) and their immediate cynodont predecessors.

**More information:** The paper 'The radiation of cynodonts and the ground plan of mammalian morphological diversity' by Marcello Ruta, Jennifer Botha-Brink, Steve Mitchell and Michael J. Benton is published in *Proceedings of the Royal Society B* 20131865 on 28th August, 2013.

Provided by University of Lincoln

APA citation: Not the end of the world: Why Earth's greatest mass extinction was the making of modern mammals (2013, August 28) retrieved 11 May 2021 from <https://phys.org/news/2013-08-world-earth-greatest-mass-extinction.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*