

Reflecting on Earth's albedo

September 6 2013



Albedo – or reflecting power of a surface – over the UK and Ireland from July 2005. This image was created from processed satellite data acquired every eight days over the course of the year. Credit: ESA/UCL/GlobAlbedo

The amount of sunlight being absorbed or reflected by Earth is one of the driving forces for weather and climate. Satellites are providing this information with unprecedented accuracy.

The reflecting power of a surface is known as 'albedo'. Bright snow and ice have a high albedo, meaning they reflect [solar radiation](#) back into space, while green areas like forests and fields have a much lower albedo.


The lower the albedo, the more energy from the Sun is absorbed.

Changes in Earth's surfaces can therefore affect how much of the Sun's energy is absorbed – such as a decrease in snow cover or an increase in the area used for agriculture. If the amount of energy absorbed changes, this has an effect on Earth's energy budget and ultimately affects our weather and climate.

To help scientists build better simulations of weather and climate, ESA's GlobAlbedo project is using satellite data to map changes in Earth's reflectivity.

Led by University College London, the team used readings from the Envisat and Spot-Vegetation satellites to produce global surface albedo maps from 1998 to 2011. The maps, [available for free online](#), provide the most accurate measure of Earth's reflectivity to date.

"GlobAlbedo is the first gap-free, 1 km-resolution map of Earth's land surface with an uncertainty estimate for every pixel. This could only have been produced from satellite data," said Professor Jan-Peter Muller of University College London, leader of the GlobAlbedo project.



The ‘V’ in its name stands for Vegetation: Proba-V will fly a reduced-mass version of the Vegetation instrument currently on board the Spot satellites to provide a daily overview of global vegetation growth. The Proba satellites are part of ESA’s In-orbit Technology Demonstration Programme: missions dedicated to flying innovative technologies. Credit: ESA-P.Carril

By combining data from different satellite sensors, scientists have maximised the coverage and created a time series that can be extended to include historical as well as future [satellite measurements](#).

The maps have proven useful to a variety of users, including the UK Met Office. Scientists there have been using them to update the land surface albedo information in the Met Office's operational Global Atmosphere [weather model](#), resulting in more accurate weather predictions and climate forecasts.

"Tests show that they help to give more accurate temperature forecasts over the United States and Asia, especially in summer," said Dr Malcolm Brooks from the Met Office. "We expect to be producing operational forecasts using these data in the spring of 2014."

Other case-studies have been looking into different uses of land albedo. These include investigating the effect of anthropogenic changes in land cover on Earth's energy balance, studying how agricultural practices influence heat waves and verifying climate models.

The mapping and monitoring of Earth's albedo will continue with ESA's recently launched Proba-V satellite and the future Sentinel-3 mission being developed under Europe's Global Monitoring for Environment and Security programme, Copernicus.

Provided by European Space Agency

Citation: Reflecting on Earth's albedo (2013, September 6) retrieved 20 September 2024 from <https://phys.org/news/2013-09-earth-albedo.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.