

'Stadium waves' could explain lull in global warming

10 October 2013



This is an image of Dr. Judith Curry, chair of the School of Earth and Atmospheric Sciences at the Georgia Institute of Technology. Credit: Georgia Institute of Technology

One of the most controversial issues emerging from the recent Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5) is the failure of global climate models to predict a hiatus in warming of global surface temperatures since 1998. Several ideas have been put forward to explain this hiatus, including what the IPCC refers to as 'unpredictable climate variability' that is associated with large-scale circulation regimes in the atmosphere and ocean. The most familiar of these regimes is El Niño/La Niña, which are parts of an oscillation in the ocean-atmosphere system. On longer multi-decadal time scales, there is a network of atmospheric and oceanic circulation regimes, including the Atlantic Multidecadal Oscillation and the Pacific Decadal Oscillation.

A new paper published in the journal *Climate Dynamics* suggests that this 'unpredictable [climate](#) variability' behaves in a more predictable way than previously assumed. The paper's authors, Marcia

Wyatt and Judith Curry, point to the so-called 'stadium-wave' signal that propagates like the cheer at sporting events whereby sections of sports fans seated in a stadium stand and sit as a 'wave' propagates through the audience. In like manner, the 'stadium wave' climate signal propagates across the Northern Hemisphere through a network of ocean, ice, and atmospheric circulation regimes that self-organize into a collective tempo.

The stadium wave hypothesis provides a plausible explanation for the hiatus in warming and helps explain why climate models did not predict this hiatus. Further, the new hypothesis suggests how long the hiatus might last.

Building upon Wyatt's Ph.D. thesis at the University of Colorado, Wyatt and Curry identified two key ingredients to the propagation and maintenance of this stadium wave signal: the Atlantic Multidecadal Oscillation (AMO) and [sea ice](#) extent in the Eurasian Arctic shelf seas. The AMO sets the signal's tempo, while the sea ice bridges communication between ocean and atmosphere. The oscillatory nature of the signal can be thought of in terms of 'braking,' in which positive and negative feedbacks interact to support reversals of the circulation regimes. As a result, climate regimes—multiple-decade intervals of warming or cooling—evolve in a spatially and temporally ordered manner. While not strictly periodic in occurrence, their repetition is regular—the order of quasi-oscillatory events remains consistent. Wyatt's thesis found that the stadium wave signal has existed for at least 300 years.

The new study analyzed indices derived from atmospheric, oceanic and sea ice data since 1900. The linear trend was removed from all indices to focus only the multi-decadal component of natural variability. A multivariate statistical technique called Multi-channel Singular Spectrum Analysis (MSSA) was used to identify patterns of variability shared by all indices analyzed, which characterizes the

'stadium wave.' The removal of the long-term trend from the data effectively removes the response from long term climate forcing such as anthropogenic greenhouse gases.

The stadium wave periodically enhances or dampens the trend of long-term rising temperatures, which may explain the recent hiatus in rising [global surface temperatures](#).

"The stadium wave signal predicts that the current pause in global warming could extend into the 2030s," said Wyatt, an independent scientist after having earned her Ph.D. from the University of Colorado in 2012.

Curry added, "This prediction is in contrast to the recently released IPCC AR5 Report that projects an imminent resumption of the warming, likely to be in the range of a 0.3 to 0.7 degree Celsius rise in global mean surface temperature from 2016 to 2035." Curry is the chair of the School of Earth and Atmospheric Sciences at the Georgia Institute of Technology.

Previous work done by Wyatt on the 'wave' shows the models fail to capture the stadium-wave signal. That this signal is not seen in climate model simulations may partially explain the models' inability to simulate the current stagnation in global surface temperatures.

"Current [climate models](#) are overly damped and deterministic, focusing on the impacts of external forcing rather than simulating the natural internal variability associated with nonlinear interactions of the coupled atmosphere-ocean system," Curry said.

The study also provides an explanation for seemingly incongruous climate trends, such as how sea ice can continue to decline during this period of stalled warming, and when the sea ice decline might reverse. After temperatures peaked in the late 1990s, hemispheric surface temperatures began to decrease, while the high latitudes of the North Atlantic Ocean continued to warm and Arctic sea ice extent continued to decline. According to the 'stadium wave' hypothesis, these trends mark a transition period whereby the future decades will

see the North Atlantic Ocean begin to cool and sea ice in the Eurasian Arctic region begin to rebound.

Most interpretations of the recent decline in Arctic sea ice extent have focused on the role of anthropogenic greenhouse gas forcing, with some allowance for natural variability. Declining [sea ice extent](#) over the last decade is consistent with the stadium wave signal, and the wave's continued evolution portends a reversal of this trend of declining sea ice.

"The stadium wave forecasts that sea ice will recover from its recent minimum, first in the West Eurasian Arctic, followed by recovery in the Siberian Arctic," Wyatt said. "Hence, the sea ice minimum observed in 2012, followed by an increase of sea ice in 2013, is suggestive of consistency with the timing of evolution of the stadium-wave signal."

The stadium wave holds promise in putting into perspective numerous observations of climate behavior, such as regional patterns of decadal variability in drought and hurricane activity, the researchers say, but a complete understanding of past [climate variability](#) and projections of future climate change requires integrating the stadium-wave signal with external climate forcing from the sun, volcanoes and anthropogenic forcing.

"How external forcing projects onto the stadium wave, and whether it influences signal tempo or affects timing or magnitude of regime shifts, is unknown and requires further investigation," Wyatt said. "While the results of this study appear to have implications regarding the hiatus in warming, the stadium wave signal does not support or refute anthropogenic [global warming](#). The stadium wave hypothesis seeks to explain the natural multi-decadal component of climate variability."

More information: M.G. Wyatt, et al., "Role for Eurasian Arctic shelf sea ice in a secularly varying hemispheric climate signal during the 20th century," *Climate Dynamics*, 2013. link.springer.com/article/10.1007/s00382-013-1950-2#page-1

Provided by Georgia Institute of Technology

APA citation: 'Stadium waves' could explain lull in global warming (2013, October 10) retrieved 18 April 2021 from <https://phys.org/news/2013-10-stadium-lull-global.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.