


Test magnet reaches 13.5 tesla – a new CERN record

18 November 2013, by Sarah Charley



A niobium-tin based magnet assembly forms part of the Short Model Coil project at CERN. Credit: Juan Carlos Perez

The Short Model Coil (SMC) programme tests new magnet technologies with magnets about 30 centimetres long. The technology developed in the SMC will eventually help engineers build more powerful magnets for the Large Hadron Collider (LHC) and future accelerators.

Currently, the LHC uses niobium-titanium superconducting magnets to both bend and focus proton beams as they race around the LHC. But these magnets are not powerful enough to support stronger focusing and higher energies. So engineers are looking into a new [superconducting material](#), niobium tin.

"With the existing niobium-titanium technology, 8 [tesla](#) is about the maximum practical operation field," says engineer Juan Carlos Perez, who is leading the SMC project. "The [magnetic](#) field you can produce thanks to the new material is at least 50% higher."

Niobium tin is a superconducting material that can

generate a magnetic field in the range from 15-20 tesla. Although it was discovered before niobium titanium, it is not commonly used in accelerators because it is challenging to work with.

"Niobium tin must be heat treated at high temperatures – about 650 0C – to form the superconducting phase, and becomes extremely brittle after the heat treatment," says Perez. "The SMC project is developing technologies to master this material, working closely with US colleagues who are heavily invested in this technology."

Engineers working on the magnets for the high-luminosity upgrade of the LHC want to eventually reach magnetic fields exceeding 12 tesla, says Perez. These higher magnetic fields will allow significantly stronger bending and focusing strengths in the LHC dipoles and quadrupoles.

"Within the next 10 years we want to build a set of new 'final-focus' quadrupoles close to the LHC experiments, with higher strength, resulting in smaller beams at the LHC collision points" says Perez. "This will increase the number of collisions per second and generate more data for the experiments. In the longer term – over the next 20 years or so – niobium tin will be a key technology. It could allow engineers to increase the energy in a future circular collider by a factor five to ten times the present record at the LHC."

The present world record for niobium-tin magnets in dipole configuration is 16.1 tesla, held by an American research group at the Lawrence Berkeley National Laboratory. The most recent CERN-built SMC, using a cable with a geometry very close to that of the 11 tesla dipoles presently under development, reached 13.5 tesla. "We still have a long way to go," says Perez. "But the SMC project is a first and encouraging step in the right direction."

More information:

cds.cern.ch/record/1450907?ln=en

Provided by CERN

APA citation: Test magnet reaches 13.5 tesla – a new CERN record (2013, November 18) retrieved 27 September 2021 from <https://phys.org/news/2013-11-magnet-tesla-cern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.