


Cooperative SO₂ and NO_x aerosol formation in haze pollution

29 January 2014



This shows a) particle number size distributions and b) fractional contributions of organics, nitrate, sulfate, ammonium, and chloride to the PM₁ mass during the haze episode from 06:00 to 15:00 LT, Jan. 12, 2013. Credit: Science China Press

Air pollution in China has exhibited noticeable changes over the past 30 years, shifting from point-source pollution (around factories and industrial plants) in the 1980s to urban pollution in the 1990s. Since the start of this century, air pollution has become increasingly regional and more complex. Recent research has indicated that the cooperative transition of SO₂ and NO_x into secondary aerosols (sulfate and nitrate) played a critical role in the haze pollution episode in China in January 2013.

The coexistence of high concentrations of primary and secondary gaseous and particulate pollutants results in numerous heterogeneous reactions occurring on the surfaces of fine particles. These reactions change the oxidizing capacity of the atmosphere, chemical compositions, and the physicochemical and optical properties of the particulate matter. The overall effect is that [air pollution](#) and haze formation is accelerated. It is, therefore, important to explore the formation mechanisms of secondary aerosols during [air pollution](#) episodes. The paper, "Mechanism for the formation of the January 2013 heavy haze pollution episode over central and eastern China",

Science China: Earth Sciences, No. 1, 2014, by Professor Wang and co-workers shows that the formation of secondary sulfate aerosols from SO₂ increases in the presence of NO_x.

This study explores the cooperative transition of SO₂ and NO_x into secondary aerosols on the surfaces of carbon-containing particles through heterogeneous reactions. The formation of sulfates from SO₂ is promoted by the existence of NO_x. Results show that as the particle size increases, the fractional contributions of secondary inorganic ions, such as sulfate and nitrate, also increase (Figure 1). The hygroscopicity of the particles increase and the increased water content can accelerate the gas-liquid-solid reactions of SO₂ and NO_x, which further increases the hygroscopicity of the particles. These processes form a positive feedback mechanism that enhances the conversion of primary gas pollutants into secondary aerosols. Consequently, it is important to reduce emissions of the precursor gases of PM_{2.5} to reduce the overall PM_{2.5} concentrations in the atmosphere. The authors conclude that in central and eastern China, SO₂ and NO_x should be controlled synchronously to reduce PM_{2.5} concentrations.

More information: WANG, Y. S., YAO, L., WANG, L. L., LIU, Z. R., JI, D. S., TANG, G. Q., ZHANG, J. K., HU, B., XIN, J. Y. Mechanism for the formation of the January 2013 heavy haze pollution episode over central and eastern China. *SCIENCE CHINA Earth Sciences*, 2014, DOI: [10.1007/s11430-013-4773-4](https://doi.org/10.1007/s11430-013-4773-4) . link.springer.com/article/10.1007%2Fs11430-013-4773-4

Provided by Science China Press

APA citation: Cooperative SO₂ and NO_x aerosol formation in haze pollution (2014, January 29) retrieved 17 April 2021 from <https://phys.org/news/2014-01-cooperative-so2-nox-aerosol-formation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.