

Hubble monitors supernova in nearby galaxy M82

26 February 2014




The Hubble data are expected to help astronomers refine distance measurements to Type Ia supernovae. In addition, the observations could yield insights into what kind of stars were involved in the explosion. Hubble's ultraviolet-light sensitivity will allow astronomers to probe the environment around the site of the [supernova explosion](#) and in the interstellar medium of the host galaxy.

Because of their consistent peak brightness, Type Ia supernovae are among the best tools to measure distances in the universe. They were fundamental to the 1998 discovery of the mysterious acceleration of the expanding universe. A hypothesized repulsive force, called dark energy, is thought to cause the acceleration.

This is a Hubble Space Telescope composite image of a supernova explosion designated SN 2014J in the galaxy M82. The Jan. 31 image, shown here as an inset, was taken in visible light with Hubble's Wide Field Camera 3. This image was superimposed into a photo mosaic of the entire galaxy taken in 2006 taken with Hubble's Advanced Camera for Surveys. Credit: NASA, ESA, A. Goobar (Stockholm University), and the Hubble Heritage Team (STScI/AURA)

This is a Hubble Space Telescope composite image of a supernova explosion designated SN 2014J in the galaxy M82. At a distance of approximately 11.5 million light-years from Earth it is the closest supernova of its type discovered in the past few decades. The explosion is categorized as a Type Ia supernova, which is theorized to be triggered in binary systems consisting of a white dwarf and another star—which could be a second white dwarf, a star like our sun, or a giant star.

Astronomers using a ground-based telescope discovered the explosion on January 21, 2014. This Hubble photograph was taken on January 31, as the supernova approached its peak brightness.



The M82 galaxy before (top) and after (bottom) its new supernova on Jan. 22. Credit: UCL/University of London Observatory/Steve Fossey/Ben Cooke/Guy

Pollack/Matthew Wilde/Thomas Wright

The January 31 image, shown here as an inset, was taken in visible light with Hubble's Wide Field Camera 3. This image was superimposed into a photo mosaic of the entire galaxy taken in 2006 taken with Hubble's Advanced Camera for Surveys.

Among the other major NASA space-based observatories used in the M82 viewing campaign are Spitzer Space Telescope, Chandra X-ray Observatory, Nuclear Spectroscopic Telescope Array (NuSTAR), Fermi Gamma-ray Space Telescope, Swift Gamma Ray Burst Explorer, and the Stratospheric Observatory for Infrared Astronomy (SOFIA).

Provided by NASA's Goddard Space Flight Center

APA citation: Hubble monitors supernova in nearby galaxy M82 (2014, February 26) retrieved 20 September 2021 from <https://phys.org/news/2014-02-hubble-supernova-nearby-galaxy-m82.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.