

Southern Ocean iron cycle gives new insight into climate change

March 16 2014



An international team of researchers analyzed the available data taken from all previous studies of the Southern Ocean, together with satellite images taken of the area, to quantify the amount of iron supplied to the surface waters of the Southern Ocean.

They found that deep winter mixing, a seasonal process which carries colder and deeper, nutrient-rich water to the surface, plays the most important role in transporting iron to the surface. The iron is then able to stimulate phytoplankton growth which supports the ocean's carbon cycle and the [aquatic food chain](#)

They were also able to determine that following the winter iron surge, a recycling process is necessary to support [biological activity](#) during the

spring and summer seasons.

Oceanographer, Dr Alessandro Tagliabue, from the University's School of Environmental Sciences, said: "We combined all available iron data, matched them with physical data from autonomous profiling floats and used the latest satellite estimates of biological iron demand to explore how iron is supplied to the phytoplankton in the Southern Ocean.

"This is important because iron limits biological productivity and air to sea CO₂ exchange in this region. We found unique aspects to the iron cycle and how it is supplied by [physical processes](#), making it distinct to other nutrients.

"This means that the Southern Ocean's nutrient supply would be affected by changes to the climate system (such as winds and freshwater input) differently to other areas of the ocean.

"We need to understand these unique aspects so that they can be used to better inform global climate predictions."

Dr Jean-Baptiste Sallée, from the Centre National de la Recherche Scientifique and the British Antarctic Survey, said: "We are really excited to make this discovery because until now we didn't know the physical processes allowing iron to reach the [ocean surface](#) and maintain biological activity. The combination of strong winds and intense heat loss in winter strongly mixes the ocean surface and the mixing reaches deep [iron](#) reservoir."

The Southern Ocean comprises the southernmost waters of the world oceans that encircle Antarctica. Researchers have long known the region is crucial in the uptake of atmospheric CO₂ and that biological processes in the Southern Ocean influence the global ocean system via northward flowing currents.

The study is published in *Nature Geoscience*.

More information: Surface-water iron supplies in the Southern Ocean sustained by deep winter mixing, [DOI: 10.1038/ngeo2101](https://doi.org/10.1038/ngeo2101)

Provided by University of Liverpool

Citation: Southern Ocean iron cycle gives new insight into climate change (2014, March 16)
retrieved 26 April 2024 from

<https://phys.org/news/2014-03-southern-ocean-iron-insight-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.