


Nanoparticle trapped with laser light temporarily violates the second law of thermodynamics

March 31 2014, by Mag. Veronika Schallhart



Artistic impressions of the nanoparticle in a laser trap. Credit: Iñaki Gonzalez and Jan Gieseler

Objects with sizes in the nanometer range, such as the molecular building blocks of living cells or nanotechnological devices, are

continuously exposed to random collisions with surrounding molecules. In such fluctuating environments the fundamental laws of thermodynamics that govern our macroscopic world need to be rewritten. An international team of researchers from Barcelona, Zurich and Vienna found that a nanoparticle trapped with laser light temporarily violates the famous second law of thermodynamics, something that is impossible on human time and length scale.


They report about their results in the latest issue of the prestigious scientific journal *Nature Nanotechnology*.

Surprises at the nanoscale

Watching a movie played in reverse often makes us laugh because unexpected and mysterious things seem to happen: glass shards lying on the floor slowly start to move towards each other, magically assemble and suddenly an intact glass jumps on the table where it gently gets to a halt. Or snow starts to form a water puddle in the sun, steadily growing until an entire snowman appears as if molded by an invisible hand. When we see such scenes, we immediately realize that according to our everyday experience something is out of the ordinary. Indeed, there are many processes in nature that can never be reversed. The physical [law](#) that captures this behavior is the celebrated [second law of thermodynamics](#), which posits that the entropy of a system – a measure for the disorder of a system – never decreases spontaneously, thus favoring disorder (high entropy) over order (low entropy).

However, when we zoom into the microscopic world of atoms and molecules, this law softens up and loses its absolute strictness. Indeed, at the nanoscale the second law can be fleetingly violated. On rare occasions, one may observe events that never happen on the macroscopic scale such as, for example heat transfer from cold to hot which is unheard of in our daily lives. Although on average the second law of

thermodynamics remains valid even in [nanoscale systems](#), scientists are intrigued by these rare events and are investigating the meaning of irreversibility at the nanoscale.



Artistic impressions of the nanoparticle in a laser trap. Credit: Iñaki Gonzalez and Jan Gieseler

Nanoparticles in laser traps

Recently, a team of physicists of the University of Vienna, the Institute of Photonic Sciences in Barcelona and the Swiss Federal Institute of Technology in Zürich succeeded in accurately predicting the likelihood of events transiently violating the second law of thermodynamics. They immediately put the mathematical fluctuation theorem they derived to

the test using a tiny glass sphere with a diameter of less than 100 nm levitated in a trap of [laser light](#). Their experimental set-up allowed the research team to capture the nano-sphere and hold it in place, and, furthermore, to measure its position in all three spatial directions with exquisite precision. In the trap, the nano-sphere rattles around due to collisions with surrounding gas molecules. By a clever manipulation of the laser trap the scientists cooled the nano-sphere below the temperature of the surrounding gas and, thereby, put it into a non-equilibrium state. They then turned off the cooling and watched the particle relaxing to the higher temperature through energy transfer from the gas molecules. The researchers observed that the tiny glass sphere sometimes, although rarely, does not behave as one would expect according to the second law: the nano-sphere effectively releases heat to the hotter surroundings rather than absorbing the heat. The theory derived by the researchers to analyze the experiment confirms the emerging picture on the limitations of the second law on the nanoscale.

Nanomachines out of equilibrium

The experimental and theoretical framework presented by the international research team in the renowned scientific journal *Nature Nanotechnology* has a wide range of applications. Objects with sizes in the [nanometer range](#), such as the molecular [building blocks](#) of [living cells](#) or nanotechnological devices, are continuously exposed to a random buffeting due to the thermal motion of the molecules around them. As miniaturization proceeds to smaller and smaller scales nanomachines will experience increasingly random conditions. Further studies will be carried out to illuminate the fundamental physics of nanoscale systems out of equilibrium. The planned research will be fundamental to help us understand how nanomachines perform under these fluctuating conditions.

More information: "Dynamic Relaxation of a Levitated Nanoparticle

from a Non-Equilibrium Steady State." Jan Gieseler, Romain Quidant, Christoph Dellago, and Lukas Novotny. *Nature Nanotechnology* AOP, February 28, 2014. [DOI: 10.1038/NNANO.2014.40](https://doi.org/10.1038/NNANO.2014.40)

Provided by University of Vienna

Citation: Nanoparticle trapped with laser light temporarily violates the second law of thermodynamics (2014, March 31) retrieved 22 September 2024 from <https://phys.org/news/2014-03-nanoparticle-laser-temporarily-violates-law.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.