

# Previously unknown atmospheric phenomenon discovered

4 April 2014



In tropical thunderstorms over the West Pacific air masses and the chemical substances they contain are quickly hurled upward to the edge of the stratosphere. If there are sufficient OH molecules in the atmosphere, the air is extensively cleaned by chemical transformation processes. Where OH concentrations are low, such as those now found in large sections of the tropical West Pacific, the cleaning capacity of the atmosphere is reduced. Credit: © Markus Rex, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research

Recent research results show that an atmospheric hole over the tropical West Pacific is reinforcing ozone depletion in the polar regions and could have a significant influence on the climate of the Earth.


An international team of researchers headed by Potsdam scientist Dr. Markus Rex from the German Alfred Wegener Institute has discovered a previously unknown atmospheric phenomenon over the South Seas. Over the tropical West Pacific there is a natural, invisible hole extending over several thousand kilometres in a layer that prevents transport of most of the natural and manmade substances into the [stratosphere](#) by virtue of its chemical composition. Like in a giant

elevator, many [chemical compounds](#) emitted at the ground pass thus unfiltered through this so-called "detergent layer" of the atmosphere. Scientists call it the "OH shield". The newly discovered phenomenon over the South Seas boosts ozone depletion in the [polar regions](#) and could have a significant influence on the future climate of the Earth – also because of rising air pollution in South East Asia.

At first Dr. Markus Rex suspected a series of flawed measurements. In October 2009 the atmospheric physicist from the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI) was on board the German research vessel "Sonne" to measure trace substances in the atmosphere in the tropical West Pacific. Tried and tested a thousand times over, the ozone probes he sent up into the tropical sky with a research balloon every 400 kilometres reported – nothing. Or to be more accurate: almost nothing. The ozone concentrations in his measurements remained nearly constantly below the detection limit of approx. 10 ppbv\* in the entire vertical range from the surface of the Earth to an altitude of around 15 kilometres. Normally ozone concentrations in this part of the atmosphere are three to ten times higher.

Although low values at an altitude of around 15 kilometres were known from earlier measurements in the peripheral area of the tropical West Pacific, the complete absence of ozone at all heights was surprising. However, after a short period of doubt and various tests of the instruments it dawned on the worldwide recognised ozone specialist that he might be onto a phenomenon yet unknown to science. A few research years later and after the involvement of other colleagues came confirmation: Markus Rex and his team on board the "Sonne" had tracked down a giant natural hole over the tropical South Seas, situated in a special layer of the lower atmosphere known as the "OH shield". The research results on the newly discovered OH

minimum will be published soon in the journal "Atmospheric Chemistry and Physics", with the Institute of Environmental Physics of the University of Bremen and other international research institutions as partners.




Nearly all chemical substances produced by people, animals, plants, algae or microorganisms on the ground or in the oceans react quickly with OH and break down in this process. During this chemical self-cleaning process substances that are not easily water-soluble are transformed into water-soluble products and then washed out by precipitation. Through this mechanism OH molecules remove most substances from the atmosphere. The OH molecule is therefore also called the detergent of the atmosphere. Only extremely long-lived chemical compounds, such as methane or CFCs, also known as "ozone killers", can rise through the OH shield into the stratosphere. Credit: Markus Rex, Alfred Wegener Institute

"Even though the sky appears to be an extensively uniform space for most people, it is composed of chemically and physically very different layers," Markus Rex explains the complex makeup of the atmosphere. The air layers near the ground contain hundreds or even thousands of chemical compounds. This is why winter and spring, mountains and sea, city and forests all have a distinct smell. The great majority of these substances are broken down into water-soluble compounds in the lower kilometres of the atmosphere and are subsequently washed out by

rain. Since these processes require the presence of a certain chemical substance, the so called hydroxyl (=OH) radical, this part of the atmosphere is called the "OH shield". It acts like a huge atmospheric washing machine in which OH is the detergent.

The OH shield is part of the troposphere, as the lower part of the atmosphere is called. "Only a few, extremely long-lived compounds manage to make their way through the OH shield," says Rex, "then they also get through the tropopause and enter the stratosphere." Tropopause refers to the boundary layer between the troposphere and the next atmospheric layer above it, the stratosphere. Particularly substances that enter the stratosphere unfold a global impact. The reason for this is that once they have reached the stratosphere, their degradation products remain up there for many years and spread over the entire globe.

Extremely long-lived chemical compounds find their way to the stratosphere, even where the OH shield is intact. These include methane, nitrous oxide ("laughing gas"), halons, methyl bromide and chlorofluorocarbons (CFCs), which are notorious as "ozone killers" because they play a major role in ozone depletion in the polar regions.



Location and extent of low ozone concentrations and thus of the OH hole over the West Pacific. Fig. (a) shows the region of origin of the air in the stratosphere, Fig. (b) ozone sonde measurements (dots) and satellite measurements (coloured map) of the total amount of

ozone in the tropospheric column of air and Fig. (c) the total amount of OH in the tropospheric column of air calculated with a model. Credit: Markus Rex, Alfred Wegener Institute.


intercepted to a larger extent in the OH shield elsewhere on the globe. One example is sulphur dioxide, which has a significant impact on the climate.

After many years of research scientists now understand the complicated process of [stratospheric ozone depletion](#) very well. "Nevertheless measured ozone depletion rates were often quite a bit larger than theoretically calculated in our models," Markus Rex points out a long unsolved problem of atmospheric research. "Through the discovery of the OH hole over the tropical West Pacific we have now presumably made a contribution to solving this puzzle." And at the same time discovered a phenomenon that raises a number of new questions for climate policy. Researchers are now tackling these questions in a new research project funded by the EU with around 9 million euros, i.e. "StratoClim", which is coordinated by the Alfred Wegener Institute. Within this project a new monitoring station will be established in the tropical Westpacific, together with the Institute of Environmental Physics at the University of Bremen, Germany.

"We have to realise," reminds the Potsdam atmospheric physicist, "that chemical compounds which enter the stratosphere always have a global impact." Thanks to the OH hole that the researchers discovered over the tropical Pacific, greater amounts of brominated hydrocarbons can reach the stratosphere than in other parts of the world. Although their ascent takes place over the tropical West Pacific, these compounds amplify ozone depletion in the polar regions. Since scientists identified this phenomenon and took it into account in the modelling of stratospheric ozone depletion, their models have corresponded excellently with the actually measured data.

However, it is not only brominated hydrocarbons that enter the stratosphere over the tropical West Pacific. "You can imagine this region as a giant elevator to the stratosphere," states Markus Rex using an apt comparison. Other substances, too, rise here to a yet unknown extent while they are

**Air mass elevator into the Stratosphere**



This is how air reaches the stratosphere. Through the rapid upward transport in tropical thunderstorms they reach an area of slow large-scale ascent and rise from there through the tropopause into the stratosphere over the course of weeks. This process is most pronounced during northern hemispheric winter. Model calculations show that, during this season, this process mainly takes place over the tropical West Pacific. Due to the formation of cirrus (= ice) clouds in the extremely cold tropical tropopause, a large portion of the water-soluble chemical substances is removed from the air and cannot reach the stratosphere. OH molecules transform water-insoluble into water-soluble compounds. Hence, if the concentration of OH molecules along the dotted transport pathways shown above is high only few chemical compounds make it into the stratosphere. Conversely, the lower the OH concentration is along the transport pathways, the more chemical compounds enter the stratosphere. Credit: Yves Nowak, Alfred Wegener Institute.

Sulphur particles in the stratosphere reflect sunlight and therefore act antagonistically to atmospheric greenhouse gases like CO<sub>2</sub>, which capture the heat of the sun on the Earth. To put it simply, whereas greenhouse gases in the atmosphere heat the globe, sulphur particles in the stratosphere have a cooling effect. "South East Asia is

developing rapidly in economic terms," Markus Rex explains a problem given little attention to date. "Contrary to most industrial nations, however, little has been invested in filter technology up to now. That is why sulphur dioxide emissions are increasing substantially in this region at present."

If one takes into account that sulphur dioxide may also reach the stratosphere via the OH hole over the tropical West Pacific, it quickly becomes obvious that the atmospheric elevator over the South Seas not only boosts [ozone depletion](#), but may influence the climate of the entire Earth. In fact, the aerosol layer in the stratosphere, which is also composed of sulphur particles, seems to have become thicker in recent years. Researchers do not know yet whether there is a connection here.

But wouldn't it be a stroke of luck if air pollutants from South East Asia were able to mitigate climate warming? "By no means," Markus Rex vigorously shakes his head. "The OH hole over the South Seas is above all further evidence of how complex climate processes are. And we are still a long way off from being in a position to assess the consequences of increased sulphur input into the stratosphere. Therefore, we should make every effort to understand the processes in the atmosphere as best we can and avoid any form of conscious or unconscious manipulation that would have an unknown outcome."

Provided by Helmholtz Association of German Research Centres

APA citation: Previously unknown atmospheric phenomenon discovered (2014, April 4) retrieved 15 September 2019 from <https://phys.org/news/2014-04-previously-unknown-atmospheric-phenomenon.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*