


# Beyond graphene: Controlling properties of 2D materials

April 28 2014

---



Graphene consists of a single layer of carbon atoms. Credit: Wikimedia Commons

(Phys.org) —Researchers at The University of Manchester have shown how they can control the properties of stacks of two-dimensional materials, opening up opportunities for new, previously-unimagined electronic devices.

The isolation of [graphene](#) at the University in 2004 led to the discovery of many other 2D [crystals](#). While graphene has an unrivalled set of superlatives, these crystals cover a large range of properties: from the most conductive to isolating, from transparent to optically active.

The next step is to combine several of these crystals in a 3D stack. This way, one can create 'heterostructures' with novel functionalities – capable of delivering applications as yet beyond the imagination of scientists and commercial partners.

The first examples of such heterostructures already exist: tunnelling transistors, resonant tunnelling diodes, and solar cells.

Writing in *Nature Physics*, the scientists, led by Nobel Laureate Sir Kostya Novoselov, demonstrate that layers in such stacks can interact strongly, which helps the researchers learn how to control the properties of such heterostructures.

By controlling the relative orientation between graphene and underlying boron nitride – one of the 2D materials and an excellent insulator – the team can reconstruct the crystal structure of graphene. This leads to creation of local strains in graphene and even opening of a band-gap, which might be useful for the functionality of many [electronic devices](#).

Professor Novoselov said: "Research on heterostructures is gaining momentum, and such possibilities for controlling the properties of heterostructures might become very useful for future applications."

PhD student Colin Woods, the researcher who carried out the vast majority of the work, said: "It was extremely exciting to see that the properties of graphene can change so dramatically by simply twisting the two crystals only a fraction of a degree.

"Generally, the previous model used to describe the sort of interaction which has been observed in our experiments describes only the 1-dimensional case, but even there it produces very nontrivial solutions.

"We hope that our system will push the mathematical development of

the model to two-dimensions, where even more exciting mathematics is to be expected."

**More information:** Commensurate–incommensurate transition in graphene on hexagonal boron nitride, by C. R. Woods, L. Britnell, A. Eckmann, R. S. Ma, J. C. Lu, H. M. Guo, X. Lin, G. L. Yu, Y. Cao, R. V. Gorbachev, A. V. Kretinin, J. Park, L. A. Ponomarenko, M. I. Katsnelson, Yu. N. Gornostyrev, K. Watanabe, T. Taniguchi, C. Casiraghi, H.-J. Gao, A. K. Geim and K. S. Novoselov, *Nature Physics*, 2014. [DOI: 10.1038/nphys2954](https://doi.org/10.1038/nphys2954)

Provided by University of Manchester

Citation: Beyond graphene: Controlling properties of 2D materials (2014, April 28) retrieved 20 April 2024 from <https://phys.org/news/2014-04-graphene-properties-2d-materials.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.