


Glow-in-the-dark tool lets scientists find diseased bats

May 29 2014



This image shows long-wave ultraviolet (UV) and white-light illumination of lesions associated with white-nose syndrome. Wing from dead eastern pipistrelle (*Pipistrellus subflavus*) lit from above with hand-held 51 LED 385-nm UV flashlight shows points of orange–yellow fluorescence. Credit: Nathan Ramsay, USGS

Scientists working to understand the devastating bat disease known as white-nose syndrome (WNS) now have a new, non-lethal tool to identify

bats with WNS lesions —ultraviolet, or UV, light.

If long-wave UV light is directed at the wings of [bats](#) with white-nose syndrome, it produces a distinctive orange-yellow fluorescence. This orange-yellow glow corresponds directly with microscopic [skin lesions](#) that are the current "gold standard" for diagnosing white-nose syndrome in bats.

"When we first saw this fluorescence of a bat wing in a cave, we knew we were on to something," said Greg Turner from Pennsylvania Game Commission, who has been using this technique since 2010. "It was difficult to have to euthanize bats to diagnose WNS when the disease itself was killing so many. This was a way to get a good indication of which bats were infected and take a small biopsy for testing rather than sacrifice the whole bat."

Millions of bats in the United States have died from the fungal disease called White nose syndrome which is caused by the fungus *Pseudogymnoascus (Geomyces) destructans*. White-nose syndrome was first seen in New York during the winter of 2006. Since then, the disease has spread to 25 U.S. states and 5 Canadian provinces.



Long-wave ultraviolet and white-light illumination of lesions associated with white-nose syndrome. Points of orange–yellow fluorescence (arrows) detected on a roosting *Indiana myotis* (*Myotis sodalis*) following surface illumination with a field-portable 9-watt 368-nm fluorescent UV light. Credit: Tina Cheng

A significant problem in studying WNS has been the unreliability of visual onsite inspection when checking for WNS in bats during hibernation; the only way to confirm presence of disease was to euthanize the bats and send them back to a laboratory for testing.

"Ultraviolet light was first used in 1925 to look for ringworm fungal infections in humans," said Carol Meteyer, USGS scientist and one of the lead authors on the paper. "The fact that this technique could be transferred to bats and have such remarkable precision for indicating

lesions positive for WNS invasion is very exciting."

To test the UV light's effectiveness, bats with and without white-nose syndrome in North America were tested by the U.S. Geological Survey's National Wildlife Health Center, first using UV light, then using traditional histological techniques to verify the UV light's accuracy.

In the USGS lab testing, 98.8 percent of bats with the orange-yellow fluorescence tested positive for [white-nose syndrome](#), whereas 100 percent of those that did not fluoresce tested negative for the disease. Targeted biopsies showed that pinpoint areas of fluorescence coincided with the microscopic wing lesions that are characteristic for WNS.

Researchers in the Czech Republic then tested the UV light-assisted biopsy technique in the field, using it to collect small samples from areas of bat wing that fluoresced under UV light. In this study, 95.5 percent of wing biopsies that targeted areas of fluorescence were positive for WNS lesions, while again 100 percent of bats that did not fluoresce were negative for WNS.

Combining research from two continents demonstrates that UV diagnostics might be applicable worldwide with great sensitivity and specificity in detecting WNS.

"Moreover, the technique hurts the animal minimally and bats fly away after providing data for research," said Natalia Martinkova from the Academy of Sciences of the Czech Republic. "This makes UV [fluorescence](#) an ideal tool for studying endangered species."

More information:

www.jwildlifedis.org/doi/pdf/10.7589/2014-03-058

Provided by United States Geological Survey

Citation: Glow-in-the-dark tool lets scientists find diseased bats (2014, May 29) retrieved 22 September 2024 from

<https://phys.org/news/2014-05-glow-in-the-dark-tool-scientists-diseased.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.