

Hot tropical oceans during Pliocene greenhouse warming

29 June 2014



Tropical forest in Martinique near the city of Fond St-Denis. Credit: Wikipedia


The impact of the greenhouse gas CO₂ on the Earth's temperature is well established by climate models and temperature records over the past 100 years, as well as coupled records of carbon dioxide concentration and temperature throughout Earth history. However, past temperature records have suggested that warming is largely confined to mid-to-high latitudes, especially the poles, whereas tropical temperatures appear to be relatively stable: the tropical thermostat model.

The new results, published today in *Nature Geoscience*, contradict those previous studies and indicate that tropical [sea surface temperatures](#) were warmer during the early-to-mid Pliocene, an interval spanning about 5 to 3 million years ago.

The Pliocene is of particular interest because CO₂ concentrations then were thought to have been about 400 parts per million, the highest level of the past 5 million years but a level that was reached for the first time last summer due to human activity.

The higher CO₂ levels of the Pliocene have long been associated with a warmer world, but evidence from tropical regions suggested relatively stable temperatures.

Project leader and Director of the Cabot Institute, Professor Richard Pancost said: "These results confirm what [climate models](#) have long predicted – that although greenhouse gases cause greater warming at the poles they also cause warming in the tropics. Such findings indicate that few places on Earth will be immune to global warming and that the tropics will likely experience associated climate impacts, such as increased tropical storm intensity."



'Scanning Electron Microscopy (SEM) image of the fossilised shell of planktic foraminifer *Globigerinoides ruber*. Planktic foraminifera are a highly abundant group of unicellular calcifiers that live in surface or near-surface

waters of the open ocean. Credit: Richard Abell

Provided by University of Bristol

The scientists focussed their attention on the South China Sea which is at the fringe of a vast warm body of water, the West Pacific Warm Pool (WPWP). Some of the most useful temperature proxies are insensitive to temperature change in the heart of the WPWP, which is already at the maximum temperature they can record. By focussing on the South China Sea, the researchers were able to use a combination of geochemical records to reconstruct sea surface temperature in the past.

Not all of the records agree, however, and the researchers argue that certain tools used for reconstructing past ocean temperatures should be re-evaluated.

The paper's first author, Charlotte O'Brien added: "It's challenging to reconstruct the temperatures of the ocean many millions of years ago, and each of the tools we use has its own set of limitations. That is why we have used a combination of approaches in this investigation. We have shown that two different approaches agree – but a third approach agrees only if we make some assumptions about how the magnesium and calcium content of seawater has changed over the past 5 million years. That is an assumption that now needs to be tested."

The work was funded by the UK's Natural Environment Research Council and is ongoing.

Dr Gavin Foster at the University of Southampton is particularly interested in coupling the temperature records with improved estimates of Pliocene [carbon dioxide](#) levels. He said: "Just as we continue to challenge our [temperature](#) reconstructions we must challenge the corresponding carbon dioxide estimates. Together, they will help us truly understand the natural sensitivity of the Earth system and provide a better framework for predicting future climate change."

More information: *Nature Geoscience*,
[dx.doi.org/10.1038/ngeo2194](https://doi.org/10.1038/ngeo2194)

APA citation: Hot tropical oceans during Pliocene greenhouse warming (2014, June 29) retrieved 20 June 2019 from <https://phys.org/news/2014-06-hot-tropical-oceans-pliocene-greenhouse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.