

# Citizen scientists take on problem of Pacific plastics

August 11 2014, by Lily Whiteman

---



Data related to beached debris aids efforts to reduce the impacts of marine debris on coastal wildlife. Here, COASST interns look for small debris (less than 5cm) in a plot. Credit: The Coastal Observation and Seabird Survey Team

In the classic 1967 movie, *The Graduate*, a newly minted college graduate played by Dustin Hoffman is told by an older friend that the future would be guided by "one word: plastics." Although the older

man's prediction did not refer to the future of marine ecology, the state of the world's oceans has, ironically, borne it out.

Most recently, footage of the Japanese tsunami of 2011 and the search for Malaysia Airlines Flight 370 have vividly underscored the deterioration of some of the world's most remote seas, showing them to be veritable [garbage](#) soups, awash in chunks of junk including metals, glass, rubber, medical waste, fishing gear, but most of all, plastics.

Plastic debris of all sizes, including bottles, cups, bags, toothbrushes and grain-sized pieces, represents from 60 to 80 percent of the world's total marine debris, according to the EPA.

## **Garbage in; garbage out**

Marine garbage comes from many sources, including galley waste and other trash from ships, recreational boaters, fishermen and offshore oil and gas facilities; boxes and merchandise from large containers ships; rivers that carry storm water-runoff and other waste; poorly maintained landfills; and beach litter.

Studies show that marine plastic harms 267 species worldwide, including species of [sea turtles](#), seabirds and marine mammals—some of which are endangered. Animals are most likely to be hurt by marine garbage after becoming entangled in it or by mistaking it for prey and eating it. For example, sea turtles commonly eat indigestible, potentially gut-clogging plastic objects because these objects look so much like one of their favorite foods: Jellyfish.

## **The Great Pacific Garbage Patch**

Due to varied oceanic and atmospheric forces, marine garbage is concentrated in five major gyres in the world's oceans. One of these is the North Pacific gyre—commonly called the Great Pacific Garbage Patch—which occupies most of the northern Pacific Ocean, an area of about 10 million square miles. One study suggests that the total mass of plastic in the Great Pacific Garbage Patch is six times that of plankton! Debris from the patch commonly washes up on beaches in Hawaii, along the Pacific Northwest coast and in Alaska.

## **Citizen scientists pitch in**

In light of the sheer physical enormity of the Great Pacific Garbage Patch and the complexity of its causes, what can we possibly do about it?

Perhaps help protect some vulnerable populations of wildlife from marine garbage in coastal regions, according to the Coastal Observation and Seabird Survey Team (COASST)—a citizen science group that monitors marine resources and ecosystem health at more than 350 beaches from northern California to Alaska. Although COASST, which receives funding from the National Science Foundation, has long focused on collecting data on beach-cast seabird carcasses as an indicator of coastal health, the group will soon also focus on collecting data on beached marine debris. Resulting data could be used to help support efforts to reduce the impacts of marine debris on coastal wildlife.

## **Identifying Vulnerable Beaches**

Here's how COASST's marine debris program will work: on a monthly basis, COASST volunteers will collect data on the locations and characteristics of beached marine debris. Debris data can be intersected

with information about the location and abundance of coastal wildlife. Resulting analyses can be used to help prioritize the where, when and what of beach cleanups.

For example, suppose scientists identified as a particular threat the potential for seals to become entangled in marine debris with "loops"—discarded plastic holders for six packs, packing straps, boat line and fishing nets or anything else a seal could stick its head through. Volunteers could sift through the COASST [marine debris](#) database to identify the locations of large concentrations of "loopy" debris during periods when seals are also abundant.

Provided by National Science Foundation

Citation: Citizen scientists take on problem of Pacific plastics (2014, August 11) retrieved 1 May 2024 from <https://phys.org/news/2014-08-citizen-scientists-problem-pacific-plastics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.