

# First indirect evidence of so-far undetected strange baryons

August 19 2014

---



Brookhaven theoretical physicist Swagato Mukherjee. Credit: Brookhaven National Laboratory

(Phys.org) —New supercomputing calculations provide the first evidence that particles predicted by the theory of quark-gluon interactions but never before observed are being produced in heavy-ion collisions at the Relativistic Heavy Ion Collider (RHIC), a facility that is dedicated to studying nuclear physics. These heavy strange baryons, containing at least one strange quark, still cannot be observed directly, but instead make their presence known by lowering the temperature at which other strange baryons "freeze out" from the quark-gluon plasma (QGP) discovered and created at RHIC, a U.S. Department of Energy (DOE) Office of Science user facility located at DOE's Brookhaven National Laboratory.

RHIC is one of just two places in the world where scientists can create and study a primordial soup of unbound quarks and gluons-akin to what existed in the early universe some 14 billion years ago. The research is helping to unravel how these building blocks of matter became bound into hadrons, particles composed of two or three quarks held together by gluons, the carriers of nature's strongest force.

"Baryons, which are hadrons made of three quarks, make up almost all the matter we see in the universe today," said Brookhaven theoretical physicist Swagato Mukherjee, a co-author on a paper describing the new results in *Physical Review Letters*. "The theory that tells us how this matter forms-including the protons and neutrons that make up the nuclei of atoms-also predicts the existence of many different baryons, including some that are very heavy and short-lived, containing one or more heavy 'strange' quarks. Now we have indirect evidence from our calculations and comparisons with experimental data at RHIC that these predicted higher mass states of strange baryons do exist," he said.

Added Berndt Mueller, Associate Laboratory Director for Nuclear and Particle Physics at Brookhaven, "This finding is particularly remarkable because strange quarks were one of the early signatures of the formation

of the primordial [quark-gluon plasma](#). Now we're using this QGP signature as a tool to discover previously unknown baryons that emerge from the QGP and could not be produced otherwise."

## **Freezing point depression and supercomputing calculations**

The evidence comes from an effect on the thermodynamic properties of the matter nuclear physicists can detect coming out of collisions at RHIC. Specifically, the scientists observe certain more-common strange baryons (omega baryons, cascade baryons, lambda baryons) "freezing out" of RHIC's quark-gluon plasma at a lower temperature than would be expected if the predicted extra-heavy strange baryons didn't exist.

"It's similar to the way table salt lowers the freezing point of liquid water," said Mukherjee. "These 'invisible' hadrons are like salt molecules floating around in the hot gas of hadrons, making other particles freeze out at a lower temperature than they would if the 'salt' wasn't there."

To see the evidence, the scientists performed calculations using lattice QCD, a technique that uses points on an imaginary four-dimensional lattice (three spatial dimensions plus time) to represent the positions of quarks and gluons, and complex mathematical equations to calculate interactions among them, as described by the theory of quantum chromodynamics (QCD).

"The calculations tell you where you have bound or unbound quarks, depending on the temperature," Mukherjee said.

The scientists were specifically looking for fluctuations of conserved baryon number and strangeness and exploring how the calculations fit with the observed RHIC measurements at a wide range of energies.

The calculations show that inclusion of the predicted but "experimentally uncharted" strange baryons fit better with the data, providing the first evidence that these so-far unobserved particles exist and exert their effect on the freeze-out temperature of the observable particles.

These findings are helping physicists quantitatively plot the points on the phase diagram that maps out the different phases of nuclear matter, including hadrons and quark-gluon plasma, and the transitions between them under various conditions of temperature and density.

"To accurately plot points on the phase diagram, you have to know what the contents are on the bound-state, hadron side of the transition line—even if you haven't seen them," Mukherjee said. "We've found that the higher mass states of strange baryons affect the production of ground states that we can observe. And the line where we see the ordinary matter moves to a lower temperature because of the multitude of higher states that we can't see."

**More information:** "Additional Strange Hadrons from QCD Thermodynamics and Strangeness Freezeout in Heavy Ion Collisions" [journals.aps.org/prl/abstract/ ... ysRevLett.113.072001](https://journals.aps.org/prl/abstract/...ysRevLett.113.072001)

Provided by Brookhaven National Laboratory

Citation: First indirect evidence of so-far undetected strange baryons (2014, August 19) retrieved 26 April 2024 from <https://phys.org/news/2014-08-indirect-evidence-so-far-undetected-strange.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.