


Plant life forms in the fossil record: When did the first canopy flowers appear?

September 2 2014



Photograph of the research canopy crane in the Parque Natural San Lorenzo (tropical rainforest), Smithsonian Tropical Research Institute, Panamá, and a detail of the venation of a fossil angiosperm leaf from the Paleocene (Cerrejón formation, Colombia). Leaf vein density can be used in the fossil record to shed light on the origin of the Neotropical forest. Designed by Lina Gonzalez, Smithsonian Tropical Research Institute (Panamá). Photos by Smithsonian Tropical Research Institute, and Andrés Baresch, Stanford University.

Most plant fossils are isolated organs, making it difficult to reconstruct the type of plant life or its ecosystem structure. In their study for *GEOLOGY*, published online on 28 Aug. 2014, researchers Camilla Crifò and colleagues used leaf vein density, a trait visible on leaf compression fossils, to document the occurrence of stratified forests with a canopy dominated by flowering plants.

Using a 40-meter-tall canopy crane equipped with a gondola, they were able to collect leaves from the very top of trees in Panama and the United States. They measured leaf vein density in 132 species from two Panamanian tropical forests and one temperate forest in Maryland (USA). The team also compared the leaf vein values of canopy-top and forest-bottom leaves (i.e., [leaf litter](#) on the forest floor).

The authors show that venation density, like plant metabolism (i.e., transpiration and photosynthesis), is higher in the leaves located in the [forest canopy](#) and decreases in leaves at lower levels. Furthermore, they found that leaves from the [forest floor](#), which are the closest analog to fossil floras, preserve this pattern.

The team also reanalyzed vein density data from the literature from the Early Cretaceous (132.5 million years ago) to the Paleocene (58 million years ago) to determine when flowering plants became part of the upper forest canopy. Vein density values similar to present ones appeared about 58 million years ago, indicating that the emergence of [flowering plants](#) in the canopy occurred by the Paleocene.

More information: Variations in angiosperm leaf vein density have implications for interpreting life form in the fossil record, Camilla Crifò et al., Published online on 28 Aug. 2014; [dx.doi.org/10.1130/G35828.1](https://doi.org/10.1130/G35828.1)

Provided by Geological Society of America

Citation: Plant life forms in the fossil record: When did the first canopy flowers appear? (2014, September 2) retrieved 20 September 2024 from <https://phys.org/news/2014-09-life-fossil-canopy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.