

Icelandic volcano system has been spewing lava since early September

October 29 2014



An overflight view taken on Sept. 8 shows the main eruptive site, known as "Baugur" (Icelandic for "ring" or "circle"). The eruption is the result of the spreading apart of two tectonic plates, the Eurasian and North American.

Jeffrey Karson, a Syracuse University geologist who recently traveled to Iceland to monitor the early stages of the eruption, says the lava field now covers more than 22 square miles (or 14,000 acres), nearly the size of Manhattan.

Iceland, which is made up of [lava flows](#), hasn't witnessed anything of the sort in 40 years. The country's most recent major [eruption](#) took place in 2010, when ash from the Eyafjallajökull volcano disrupted air travel for more than a week across parts of Europe.

"The lava that erupted at Eyafjallajökull was much less extensive than what we're seeing now," says Karson, an Earth sciences professor in the College of Arts and Sciences. "This one doesn't show any signs of slowing down.

An expert in structural geology and tectonics and the co-founder of the Syracuse University Lava Project, Karson regularly travels to Iceland to study faulting and volcanic structures. His latest trip in September gave him a first-hand glimpse—much of it from close range—of the ongoing eruption fed by the Bárðarbunga volcano, located under the country's most extensive glacier. On average, the system produces enough [lava](#), every five minutes, to fill an area the size of a football field.

"The eruption is the result of the spreading apart of two tectonic plates [the Eurasian Plate and North American Plate], which are literally pulling Iceland apart," says Karson, who is collaborating with investigators at the University of Iceland's Institute of Earth Sciences. "As the plates diverge, magma from deep in the Earth is injected upward to fill the gaps. Individual eruptions can last for months, or even years."

Karson adds that the eruption has been one of the most closely studied ever: "Iceland is a natural laboratory that allows us to study volcanoes and faulting across a range of disciplines, including geochemistry, geophysics, geology and petrology. Everything we learn there adds to our understanding of how the Earth works."



A view of the eruptive plume, captured Sept. 9, was taken from about 50 kilometers to the north of the main vents. Jeffrey Karson, Earth sciences professor in the College of Arts and Sciences, describes this eruption as one of the most closely studied ever, with Iceland as a natural laboratory.



Jeffrey Karson, an expert in structural geology and tectonics, gets within two kilometers of the main eruptive vents. His collaboration with investigators at the University of Iceland Institute of Earth Sciences gives him a first-hand view of the early stages of the eruption.



Open fissures above subsurface magma conduit or channels feed the volcano's eruptive vents.



This evening view of a lava fountain at the volcano's central vent shows striking contrast between molten lava and surrounding rock and glacial ice. Erupting lava is thrown 50-100 meters into the air above the vents.

Provided by Syracuse University

Citation: Icelandic volcano system has been spewing lava since early September (2014, October 29) retrieved 19 September 2024 from <https://phys.org/news/2014-10-icelandic-volcano-spewing-lava-early.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.