

West Antarctic melt rate has tripled

December 2 2014



UCI and NASA glaciologists, including Isabella Velicogna and Tyler Sutterley, have discovered that the melt rate of glaciers in West Antarctica has tripled, with the loss of a Mt. Everest's worth of water weight every two years.

A comprehensive, 21-year analysis of the fastest-melting region of Antarctica has found that the melt rate of glaciers there has tripled during the last decade.

The glaciers in the Amundsen Sea Embayment in West Antarctica are hemorrhaging ice faster than any other part of Antarctica and are the

most significant Antarctic contributors to [sea level rise](#). This study is the first to evaluate and reconcile observations from four different measurement techniques to produce an authoritative estimate of the amount and the rate of loss over the last two decades.

"The mass loss of these glaciers is increasing at an amazing rate," said scientist Isabella Velicogna, jointly of the UC Irvine and NASA's Jet Propulsion Laboratory. Velicogna is a coauthor of a paper on the results, which has been accepted for Dec. 5 publication in the journal *Geophysical Research Letters*.

Lead author Tyler Sutterley, a UCI doctoral candidate, and his team did the analysis to verify that the melting in this part of Antarctica is shifting into high gear. "Previous studies had suggested that this region is starting to change very dramatically since the 1990s, and we wanted to see how all the different techniques compared," Sutterley said. "The remarkable agreement among the techniques gave us confidence that we are getting this right."

The researchers reconciled measurements of the mass balance of glaciers flowing into the Amundsen Sea Embayment. Mass balance is a measure of how much ice the glaciers gain and lose over time from accumulating or melting snow, discharges of ice as icebergs, and other causes. Measurements from all four techniques were available from 2003 to 2009. Combined, the four data sets span the years 1992 to 2013.

The glaciers in the embayment lost mass throughout the entire period. The researchers calculated two separate quantities: the total amount of loss, and the changes in the rate of loss.

The total amount of loss averaged 83 gigatons per year (91.5 billion U.S. tons). By comparison, Mt. Everest weighs about 161 gigatons, meaning the Antarctic [glaciers](#) lost a Mt.-Everest's-worth amount of water weight

every two years over the last 21 years.

The rate of loss accelerated an average of 6.1 gigatons (6.7 billion U.S. tons) per year since 1992.

From 2003 to 2009, when all four observational techniques overlapped, the melt rate increased an average of 16.3 gigatons per year—almost three times the rate of increase for the full 21-year period. The total amount of loss was close to the average at 84 gigatons.

The four sets of observations include NASA's Gravity Recovery and Climate Experiment (GRACE) satellites, laser altimetry from NASA's Operation IceBridge airborne campaign and earlier ICESat satellite, radar altimetry from the European Space Agency's Envisat satellite, and mass budget analyses using radars and the University of Utrecht's Regional Atmospheric Climate Model.

The scientists noted that glacier and ice sheet behavior worldwide is by far the greatest uncertainty in predicting future sea level. "We have an excellent observing network now. It's critical that we maintain this network to continue monitoring the changes," Velicogna said, "because the changes are proceeding very fast."

Provided by University of California, Irvine

Citation: West Antarctic melt rate has tripled (2014, December 2) retrieved 25 April 2024 from <https://phys.org/news/2014-12-west-antarctic-tripled.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.