


Image: Coloring the sea around the Pribilof Islands

12 January 2015



Credit: NASA/Landsat 8

The Operational Land Imager (OLI) on Landsat 8 captured this view of a phytoplankton bloom near Alaska's Pribilof Islands on Sept. 22, 2014.

The Pribilofs are surrounded by nutrient-rich waters in the Bering Sea. The milky green and light blue shading of the water indicates the presence of vast populations of microscopic phytoplankton—mostly coccolithophores, which have calcite scales that appear white in satellite images. Such phytoplankton form the foundation of a tremendously productive habitat for fish and birds.

Blooms in the Bering Sea increase significantly in springtime, after winter ice cover retreats and nutrients and freshened water are abundant near

the ocean surface.

Phytoplankton populations plummet in summertime as the water warms, surface nutrients are depleted by blooms, and the plant-like organisms are depleted by grazing fish, zooplankton, and other marine life.

By autumn, storms can stir [nutrients](#) back to the surface and cooler waters make better bloom conditions.

Provided by NASA

APA citation: Image: Coloring the sea around the Pribilof Islands (2015, January 12) retrieved 6 March 2021 from <https://phys.org/news/2015-01-image-sea-pribilof-islands.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.