


TRMM and GPM satellites measure rainfall rates in Typhoon Higos

February 10 2015, by Rob Gutro



On Feb. 9, 2015 GPM's GMI measured rain falling at a rate of over 35.5 mm (1.3 inches) southwest of the center of circulation. Credit: NASA/JAXA, Hal Pierce

The Tropical Rainfall Measuring Mission or TRMM satellite and the Global Precipitation Measurement or GPM core satellites can calculate rainfall rates occurring in a storm from their orbits in space. TRMM and GPM both saw moderate rainfall occurring in Typhoon Higos as it moved over open waters of the Northwestern Pacific Ocean.

Both TRMM and GPM are managed by NASA and the Japan Aerospace Exploration Agency called JAXA.


Both TRMM and the GPM core observatory satellite had good looks at Higos' [rainfall](#) on February 9, as the passed overhead. Rainfall rates were derived from GPM's Microwave Imager (GMI) at 0206 UTC and the TRMM Microwave Imager (TMI) three hours later at 0546 UTC. GPM's GMI measured rain falling at a rate of over 35.5 mm (1.3 inches) southwest of the center of circulation and TRMM's TMI measured rain falling at the same rate.

On Feb. 10, 2015 at 03:10 UTC (Feb. 9 at 10:10 p.m. EST) the Moderate Resolution Imaging Spectroradiometer (MODIS) instrument aboard NASA's Aqua satellite captured a visible image of Tropical Cyclone Higos as its eye was filling in with clouds.

On Feb. 10 at 1500 UTC (10 a.m. EST), Higos was still a typhoon with [maximum sustained winds](#) near 100 knots (115.1 mph/185.2 kph). It was centered near 14.7 north latitude and 154.1 east longitude, about 451 nautical miles (519 miles/835 km) north-northeast of Chuuk. Higos was moving to the north at 5 knots (5.7 mph/9.2 kph).

At that time, the Joint Typhoon Warning Center (JTWC) noted that animated enhanced infrared satellite imagery showed that Higos had started to weaken, and that the eye feature had dissipated. In addition,

deep (strong) convection (rapidly rising air that forms the thunderstorms that make up a tropical cyclone) had become less organized.




On Feb. 9, 2015 TRMM's TMI also measured rain falling at a rate of over 35.5 mm (1.3 inches) west of the center of circulation. Credit: NASA/JAXA, Hal Pierce

A couple of factors are expected to continue weakening Higos and bring it to an end in three days, according to JTWC. Decreasing ocean heat

and the possibility of dry air moving into the storm are expected to lead to Higos' demise by Feb. 13.

The JTWC forecast indicates that Higos will begin moving to the northeast and weaken while staying over the [open waters](#) of the Pacific Ocean and well northeast of Guam.



On Feb. 10, 2015 at 03:10 UTC NASA's Aqua satellite captured this visible image of Tropical Cyclone Higos as its eye was filling in with clouds. Credit: NASA Goddard MODIS Rapid Response Team

Provided by NASA's Goddard Space Flight Center

Citation: TRMM and GPM satellites measure rainfall rates in Typhoon Higos (2015, February 10) retrieved 19 September 2024 from <https://phys.org/news/2015-02-trmm-gpm-satellites-rainfall-typhoon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.