


Microfluidic diamond sensor: Moving bio particles magnetically

February 19 2015



A diamond nanocrystal (white object to the right of center) is used to map the magnetic field around a particle (red object at center). The particle floats in a shallow bath of ionic liquid. The particle can be moved about (dotted line) with great precision by making the liquid flow using voltages applied to electrodes (4 shiny rods). Inset: the NV center at the heart of the diamond nano-crystal reacts to a combination of incoming green laser light, radio-frequency waves (magenta), and the magnetism of the nearby micro-particle. If all these fields have just the right values the NV center will emit red light. The observed light provides a measure of the micro-particle's magnetic field. Credit: Kelley/JQI

Measuring faint magnetic fields is a trillion-dollar business. Gigabytes

of data, stored and quickly retrieved from chips the size of a coin, are at the heart of consumer electronics. Even higher data densities can be achieved by enhancing magnetic detection sensitivity—perhaps down to nano-tesla levels.

Greater magnetic sensitivity is also useful in many scientific areas, such as the identification of biomolecules such as DNA or viruses. This research must often take place in a warm, wet environment, where clean conditions or low temperatures are not possible. JQI scientists address this concern by developing a diamond sensor that operates in a fluid environment. The sensor makes magnetic maps (with a 17 micro-tesla sensitivity) of small particles (a stand-in for actual bio-molecules) with a spatial resolution of about 50 nm. This is probably the most sensitive magnetic measurement conducted at room temperature in microfluidics.

The results of the new experiment conducted by JQI scientist Edo Waks (a professor at the University of Maryland) and his associates appear in the journal *NanoLetters* .

Diamond NV centers

At the heart of the sensor is a tiny diamond nano-crystal. This diamond, when brought close to a magnetic particle while simultaneously being bathed in [laser light](#) and a subtle microwave signal, will fluoresce in a manner proportional to the strength of the particle's own [magnetic field](#). Thus light from the diamond is used to map magnetism.

How does the diamond work and how is the particle maneuvered near enough to the diamond to be scanned?

The diamond nanocrystal is made in the same process by which synthetic [diamonds](#) are formed, in a process called chemical vapor deposition. Some of the diamonds have tiny imperfections, including occasionally

nitrogen atoms substituting for carbon atoms. Sometimes a carbon atom is missing altogether from the otherwise tightly-coordinated diamond solid structure. In those cases where the nitrogen (N) and the vacancy (V) are next to each other, an interesting optical effect can occur. The NV combination acts as a sort of artificial atom called an NV color center. If prompted by the right kind of green laser, the NV center will shine. That is, it will absorb green laser light and emit red light, one photon at a time.

The NV emission rate can be altered in the presence of magnetic fields at the microscopic level. For this to happen, though, the internal energy levels of the NV center has to be just right, and this comes about when the center is exposed to signals from the radio-frequency source (shown at the edge of the figure) and the fields emitted by the nearby [magnetic particle](#) itself.

The particle floats in a shallow lake of de-ionized- water based solution in a setup called a microfluidic chip. The diamond is attached firmly to the bottom of this lake. The particle moves, and is steered around the chip when electrodes positioned in the channels coax ions in the liquid into forming gentle currents. Like a ship sailing to Europe with the help of the Gulf Stream, the particle rides these currents with sub-micron control. The particle can even be maneuvered in the vertical direction by an external magnetic coil (not shown in the drawing).

"We plan to use multiple diamonds in order to do complex vectorial magnetic analysis.," says graduate student Kangmook Lim, the lead author on the publication. "We will also use floating diamonds instead of stationary, which would be very useful for scanning nano- magnetism of biological samples."

More information: "Scanning localized magnetic fields in a microfluidic device with a single nitrogen vacancy center," Kangmook

Lim, Chad Ropp, Benjamin Shapiro, Jacob M. Taylor, Edo Waks, *Nano Letters*, 5 February 2015; pubs.acs.org/doi/abs/10.1021/nl503280u

Provided by Joint Quantum Institute

Citation: Microfluidic diamond sensor: Moving bio particles magnetically (2015, February 19)
retrieved 20 September 2024 from
<https://phys.org/news/2015-02-microfluidic-diamond-sensor-bio-particles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.