

# Exploding stars help to understand thunderclouds on Earth

22 April 2015



Particle shower initiated by a cosmic ray reaches LOFAR through a thundercloud. Credit: Radboud University

How is lightning initiated in thunderclouds? This is difficult to answer - how do you measure electric fields inside large, dangerously charged clouds? It was discovered, more or less by coincidence, that cosmic rays provide suitable probes to measure electric fields within thunderclouds. This surprising finding is published in *Physical Review Letters* on April 24th. The measurements were performed with the LOFAR radio telescope located in the Netherlands.

'We used to throw away LOFAR [measurements](#) taken during [thunderstorms](#). They were too messy,' says astronomer Pim Schellart. 'Well, we didn't actually throw them away of course, we just didn't analyze them.' Schellart, who completed his PhD in March this year at Radboud University in

Nijmegen and is supervised by Prof. Heino Falcke, is interested in [cosmic rays](#). These high-energy particles, originating from exploding stars and other astrophysical sources, continuously bombard Earth from space.

High in the atmosphere these particles strike atmospheric molecules and create 'showers' of [elementary particles](#). These showers can also be measured from the radio emission that is generated when their constituent particles are deflected by the magnetic field of the Earth. The radio emission also gives information about the original particles. These measurements are routinely conducted with LOFAR at ASTRON in Dwingeloo, but not during thunderstorms.

## Modeling

That changed when the data were examined in a collaborative effort with astrophysicist Gia Trinh, Prof. Olaf Scholten from the University of Groningen and lightning expert Ute Ebert from the Centrum Wiskunde & Informatica in Amsterdam.

'We modeled how the electric field in thunderstorms can explain the different measurements. This worked very well. How the [radio emission](#) changes gives us a lot of information about the electric fields in thunderstorms. We could even determine the strength of the electric field at a certain height in the cloud.' says Schellart.

This field can be as strong as 50 kV/m. This translates into a voltage of hundreds of millions of volts over a distance of multiple kilometers: a thundercloud contains enormous amounts of energy.

## Dangerous charge

Lightning is a highly unpredictable natural phenomenon that inflicts damage to infrastructure and claims victims around the world. This new

method to measure electric fields in thunderclouds will contribute to a better understanding and ultimately better predictions of lightning activity. Current measurement methods from planes, balloons or little rockets are dangerous and too localized. Most importantly the presence of the measurement equipment influences the measurements. Cosmic rays probe the thunderclouds from top to bottom. Moving at almost the speed of light they provide a near instantaneous 'picture' of the electric fields in the cloud. Moreover, they are created by nature and are freely available.

'This research is an exemplary form of interdisciplinary collaboration between astronomers, particle physicists and geophysicists', says Heino Falcke. 'We hope to develop the model further to ultimately answer the question: how is lightning initiated within thunderclouds?'

**More information:** Probing atmospheric electric fields in thunderstorms through radio emission from cosmic-ray induced air showers, P. Schellart et al. *Physical Review Letters* 24 April 2015.

Provided by University of Groningen

APA citation: Exploding stars help to understand thunderclouds on Earth (2015, April 22) retrieved 26 May 2019 from <https://phys.org/news/2015-04-stars-thunderclouds-earth.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*