

# Acid saline groundwaters and lakes of southern Western Australia

June 1 2015

---



Shallow acid saline water in Gneiss Lake, near Grass Patch in Western Australia, is an example of end-member continental brines. Orange iron oxide staining and

white halite and gypsum precipitate on Precambrian quartzite gravel in lake. Photo taken in January 2008, when lake water was undergoing evapo-concentration and had pH 2.0 and salinity 28 percent total dissolved solids. Credit: Kathleen Benison, *GSA Today* June 2015.

The "wheat belt" and "gold fields" of southern Western Australia are associated with a regional acid saline groundwater system. Groundwaters hosted in the Yilgarn Craton there have pH levels as low as 2.4 and salinities as high as 28%, which have greatly affected bedrock and subsurface sediments. This is manifested above ground as hundreds of shallow, ephemeral acid saline lakes.

In the June issue of *GSA Today*, Kathleen Benison of West Virginia University and Brenda Bowen of the University of Utah write that the limited volume of groundwater, in combination with its acidity, salinity, and high concentrations of some metals, make southern Western Australia a difficult place for human habitation.


The overarching question addressed by this study is "How did the extreme acidity form here?" The authors discuss the combination of processes that make up these shallow lakes and the groundwater that feeds them, which is not only dependent upon the host rock lithology, mineralogy, climate, weathering, organisms, and time, but also on mining and agriculture in the region.

Benison and Bowen write, "In the twentieth century both agriculture and mining had local influence on acid brine groundwater. A government-sponsored effort to turn the semi-arid eucalypt forests of inland southern Western Australia to crop and ranchland promoted the deforestation of the 'wheat belt' region. With fewer trees to soak up the acid saline groundwater, the water table rose. Ranchers realized that cattle and

sheep did not thrive with acid brines. Farmers found the only successful crops were grown high above the water table and irrigated with desalinized seawater piped a distance of hundreds of kilometers."

Mining efforts have also used desalinized seawater pipelines. Both mining and agriculture import fresher water to the [groundwater system](#) and may be responsible for changing the volume of groundwater slightly, as well as potentially causing dissolution of some subsurface halite and other chemical sediments, and, perhaps in turn, increasing [groundwater salinity](#).

Looking forward at further research areas, the authors write, "Extreme acid brine environments similar to those in southern Western Australia have been recognized on Earth and Mars. In particular, some mid-Permian continental environments hosted extremely acid saline lakes and groundwaters that deposited redbeds and evaporites. The temporal and geographic extent of these Permian acid brine settings, and their relationship to Permian climate change and the end Permian mass extinction, are open scientific questions. Understanding the origin, evolution, and maintenance of modern natural acid brine environments may lead to more informed paleoenvironmental, paleoclimatic, and paleobiological interpretations about ancient [acid](#) brines.



Basic geological units of Australia, after Addario et al., created in Arcinfo GIS from public domain geological mapping data, GFDL free use. Dark brown: Craton.

**More information:** The evolution of end-member continental waters: The origin of acidity in southern Western Australia, *GSA Today*, v. 25, no. 6, [dx.doi.org/10.1130/GSATG231A.1](https://doi.org/10.1130/GSATG231A.1).

Provided by Geological Society of America

Citation: Acid saline groundwaters and lakes of southern Western Australia (2015, June 1)  
retrieved 20 September 2024 from

<https://phys.org/news/2015-06-acid-saline-groundwaters-lakes-southern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.