


# Aircraft debris looks like it's from MH370 – now can we find the rest?

31 July 2015, by



Currents map the debris' possible route across the ocean. Michael Pidwirny

It appears that the debris washed ashore on Reunion, an island east of Madagascar, may be from the missing Malaysia Airlines Flight MH370 which disappeared in March 2014, believed lost at sea somewhere to the west of Australia.

Reunion lies 500km east of Madagascar near the island of Mauritius, around 4,000km from the area where search efforts for the missing aircraft have been concentrated. That's a huge distance to travel, even in the 500 or so days it has been since the crash. Is this possible from an oceanographic perspective?

Certainly, it's possible that aircraft debris – which is built to be relatively lightweight, otherwise it would be difficult for the aircraft to fly – can float quite close to the surface. The near-surface [ocean currents](#) in the region are mainly driven by broad wind patterns.

In the southern Indian Ocean the average long-term near surface circulation is counter clockwise, and so material that enters the ocean southwest of Australia would be carried by the West Australia current northwards, towards the equator. There it

could join the South Equatorial current moving westwards until it joined the Mozambique current travelling along the African coast and past Reunion. The distance travelled – roughly 5,000-6,000km in 15 months or so – gives an average speed of about 15cm/s. This is quite a reasonable value for the currents in the upper levels of the ocean water column where such debris might find itself suspended.

## Using ocean currents to look back into the past

Even if it is part of Flight MH370, does this provide us with any new information that could help investigators pinpoint the crash site location? Perhaps "pinpoint" is too strong a word. But it's certainly possible to use numerical ocean circulation models to trace the route the debris might have taken and suggest new regions of the ocean to search.

Such simulations come with certain assumptions. Just as a weather forecast becomes less accurate the further ahead we try to predict, attempting to re-create a journey like this through simulation becomes less certain the further back into the past we have to travel.

This is because the circulation transporting the debris would be driven by winds, and we are talking about winds over a remote part of the ocean. Consequently any uncertainty in the wind field used in the model will introduce uncertainty into the model simulations of the ocean currents, which would in turn create potentially large variability in any predictions of the location of a crash site the ocean currents dragged the [debris](#) from.

This is not to say that it's not worth doing – it may very well indicate which part of the ocean the search should focus on, and perhaps even more importantly rule out areas not worth visiting. If one piece of wreckage has emerged, then there may be more to come, which could add accuracy to the

prediction model.

Ultimately, when searching an ocean area as huge as the Indian and Southern oceans, any help is better than nothing at all.

*This story is published courtesy of [The Conversation](#) (under Creative Commons-Attribution/No derivatives).*

Source: The Conversation

APA citation: Aircraft debris looks like it's from MH370 – now can we find the rest? (2015, July 31) retrieved 19 October 2019 from <https://phys.org/news/2015-07-aircraft-debris-mh370-rest.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*