

Scientists pioneer method to track water flowing through glaciers

August 10 2015



Researchers work with seismic equipment to track meltwater running through Alaska's Yahtse Glacier. Credit: Tony Oney

Researchers for the first time have used seismic sensors to track meltwater flowing through glaciers and into the ocean, an essential step to understanding the future of the world's largest glaciers as climate changes.

The University of Texas Institute for Geophysics (UTIG) helped pioneer this new method on glaciers in Greenland and Alaska. The study will be published Aug. 10 in the journal *Geophysical Research Letters*.

Meltwater moving through a glacier into the [ocean](#) is critically important because it can increase melting and destabilize the glacier in a number of ways: The water can speed the glacier's flow downhill toward the sea; it can move rocks, boulders and other sediments toward the terminus of the glacier along its base; and it can churn and stir warm ocean water, bringing it in contact with the glacier.

"It's like when you drop an ice cube into a pot of warm water. It will eventually melt, but it will melt a lot faster if you stir that water," said Timothy Bartholomaus, a postdoctoral fellow at UTIG and the study's lead author. "Subglacial discharge provides that stirring."

The new technique offers scientists a tool for tracking [meltwater](#) at glaciers that end in the ocean, called [tidewater glaciers](#). Unlike landlocked glaciers, where scientists can simply measure the meltwater flowing in glacial rivers, there previously had not been a method available to track what's occurring within tidewater glaciers.

"All of the biggest glaciers in Greenland, all of the biggest glaciers in Antarctica, they end in the ocean," Bartholomaus said. "We need to understand how these glaciers are moving and how they are melting at their front. If we want to answer those questions, we need to know what's occurring with the meltwater being discharged from the glacier."



The brown water at the top of the picture is the subglacial discharge, or meltwater, that has flowed through Alaska's Yahtse Glacier and into the ocean. Scientists at The University of Texas at Austin have pioneered a method to track meltwater flowing through glaciers that end in the ocean. Credit: Tony Oney

UTIG research associate Jake Walter worked on the study. The team also includes researchers from the University of Alaska Southeast, the U.S. Geological Survey and the University of Alaska Fairbanks. Bartholomaus did his fieldwork while studying for his doctorate at the University of Alaska Fairbanks, but he analyzed the data and wrote the study while at UTIG.

UTIG is a research unit of The University of Texas at Austin Jackson School of Geosciences.

The team discovered the new method while trying to study earthquakes caused by iceberg calving—when large chunks of ice break off glaciers. Bartholomaeus said the ability to identify these earthquakes, known as icequakes, varied over the season, and that they were much more difficult to detect during summer because seismic [background noise](#) was obscuring the icequake signals.

The team set about trying to determine what was causing the background noise, investigating potential causes such as rainfall, iceberg calving and the movement of the glacier over the ground. Eventually, as the researchers discounted these theories, they discovered that the seismic vibrations being detected by the equipment was caused by meltwater percolating down through the glacier and weaving its way through the complicated plumbing system in the interior of the ice.

Researchers tested the theory on glaciers with meltwater rivers and found that the timing of the meltwater and the seismic signals synced perfectly. The method is very good at identifying when the glacial discharge is flowing into the ocean, Bartholomaeus said, but it will take more research to determine exactly how much water is flowing out.

"Now that we know when subglacial discharge is faster or slower, we can make better measurements of glacier change," Bartholomaeus said. "My hope is that this method will really help us understand how the glaciers and the oceans are coupled, and how the ocean might be affecting the behavior of tidewater [glaciers](#)."

More information: Subglacial discharge at tidewater glaciers revealed by seismic tremor, DOI: 10.1002/2015GL064590

Provided by University of Texas at Austin

Citation: Scientists pioneer method to track water flowing through glaciers (2015, August 10)
retrieved 20 September 2024 from
<https://phys.org/news/2015-08-scientists-method-track-glaciers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.