


# Scientists track ultrafast formation of catalyst with X-ray laser

11 August 2015



The benchmark catalyst  $\text{Fe}(\text{CO})_5$  is irradiated with ultraviolet light, causing it to lose one of its five carbon monoxide groups. The resulting reaction is probed with a soft x-ray source that provides information about chemical dynamics at an atom-specific level with femtosecond or quadrillionths of a second resolution. Credit: Helmholtz Zentrum Berlin/SciStyle, Thomas Splettstößer

At the SLAC National Accelerator Laboratory, an international team has – for the first time – precisely tracked the surprisingly rapid process by which light rearranges the outermost electrons of a metal compound known as  $\text{Fe}(\text{CO})_5$  and turns it into an active catalyst – a substance that promotes chemical reactions. The results come from using resonant inelastic x-ray scattering at SLAC, which also provides information about the chemical dynamics of the catalyst.

The new x-ray spectroscopy technique allows scientists to investigate how light rearranges the

outermost electrons of a compound in a few hundred femtoseconds, or quadrillionths of a second. Learning these details could help scientists predict and control the quick, early steps in reactions to turn sunlight and water into fuel as well as master the chemistry required to produce other renewable fuels.

The proper methods to interrogate the electronic states of transition metal complexes, such as the iron-based  $\text{Fe}(\text{CO})_5$ , during a reaction have been lacking. Recently, a team of researchers from the United States and Europe employed a technique designed to alleviate this problem. This research group used resonant inelastic x-ray scattering (RIXS) at the SLAC National Accelerator Laboratory to probe the frontier or outermost orbitals of  $\text{Fe}(\text{CO})_5$  with atomic-level and femtosecond resolution. In this work, the researchers used femtosecond duration x-ray laser pulses generated by the Linac Coherent Light Source to follow Fe-CO bond dissociation in the metal complex and the resulting catalyst formation.

The measurement is the first of its kind, showing the potential for using multi-dimensional x-ray spectroscopy to probe chemical dynamics and better understand how catalysts function. Data acquired through this technique has already addressed a long-standing question in the photochemistry of the catalyst  $\text{Fe}(\text{CO})_4$  by demonstrating that triplet and singlet reaction pathways occur in parallel even on the sub-picosecond timescale. The use of RIXS should be applicable to a variety of research fields, especially those looking to gain information on structural dynamics in ultrafast processes.

**More information:** "Orbital-specific mapping of the ligand exchange dynamics of  $\text{Fe}(\text{CO})_5$  in solution." *Nature* 520, 78-81 (2015). DOI: [10.1038/nature14296](https://doi.org/10.1038/nature14296)

Provided by US Department of Energy

APA citation: Scientists track ultrafast formation of catalyst with X-ray laser (2015, August 11) retrieved 15 June 2021 from <https://phys.org/news/2015-08-scientists-track-ultrafast-formation-catalyst.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*