

# Meat food waste has greater negative environmental impact than vegetable waste

August 14 2015

---



Approximately 31 percent of food produced in the U.S., or 133 billion pounds of food worth \$162 billion, was wasted in 2011 according to the United States Department of Agriculture (USDA). Now, University of Missouri researchers have found that the type of food wasted has a significant impact on the environment. Although less meat is wasted (on average) compared to fruits and vegetables, the researchers found that significantly more energy is used in the production of meat compared to the production of vegetables. This wasted energy is usually in the form of resources that can have negative impacts on the surrounding environment, such as diesel fuel or fertilizer being released into the environment.

"While many of us are concerned about [food waste](#), we also need to consider the resources that are wasted when we throw away edible food,"

said Christine Costello, assistant research professor at the College of Agriculture, Food and Natural Resources and co-author of the study. "Farm equipment used to feed and maintain livestock and plant and harvest crops uses a lot of [diesel fuel](#) and other utilities from fossil fuels. When people waste meat, these fuels, as well as fertilizers, are also wasted. Based on our study, we recommend that people and institutions be more conscious of not only the amount but the types of food being wasted."

During the study, pre- and post-consumer food waste was collected from four all-you-care-to-eat dining facilities over three months in 2014. Costello and her research team created a detailed inventory of the specific types of food waste: meat, vegetables or starches. The food waste also was categorized as either edible or inedible (peels and ends of [fruits and vegetables](#)).

Once the food waste was categorized, Costello and her research team analyzed greenhouse gas (GHG) emissions resulting from fertilizer use, vehicle transportation, and utility use on the farm. GHG emission estimates were measured from cradle (land preparation or animal birth) to farm gate (when the grain or animal was sent to a processing facility). Previous studies have shown that the majority of GHG emissions occur in the production stages prior to the farm products' leaving the farm.

"Based on the findings, we recommend consumers pay special attention to avoiding waste when purchasing and preparing meat; if consumers choose to prepare extra food 'just in case,' they should use plant-based foods," said co-author Ronald G. McGarvey, assistant professor at the Harry S Truman School of Public Affairs and Department of Industrial and Manufacturing Systems Engineering.

Costello and McGarvey recommend that future research should examine the effects of facilities that operate as all-you-care-to-eat versus those

that operate à la carte at the consumer level. McGarvey and Esma Birisci, a doctoral student in industrial engineering, are developing techniques to improve production as well as ordering decisions to reduce food waste and corresponding GHG emissions.

The study, "Food waste in campus dining operations: Inventory of pre- and post-consumer mass by food category, and estimation of embodied [greenhouse gas emissions](#)," was published in *Renewable Agriculture and Food Systems*.

Provided by University of Missouri-Columbia

Citation: Meat food waste has greater negative environmental impact than vegetable waste (2015, August 14) retrieved 19 September 2024 from <https://phys.org/news/2015-08-meat-food-greater-negative-environmental.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.